

INDUCIBLE ALPHAVIRAL/ORIP BASED GENE EXPRESSION SYSTEM

BACKGROUND OF THE INVENTION

5 Field of the Invention

The present invention relates to compositions and methods for the production of polypeptides and/or untranslated RNA molecules in host cells. The invention provides nucleic acid molecules, expression systems, host cells, methods and kits that are useful for the production of polypeptides and/or untranslated RNA molecules.

Related Art

Process development for biopharmaceuticals is influenced by product quality and economy of the manufacturing process. The economic production of recombinant proteins in mammalian cells is dependent on the selection of the producing cell lines. The classical approach is the use of stable expression systems. These systems are based on chromosomal integration of an expression plasmid into the genome of the host cell.

An alternative to stable expression systems is transient gene expression. For example, transient gene expression in, e.g., mammalian cells, at reactor scale is becoming increasingly important for the rapid production of recombinant proteins. Large scale applications of transient expression systems for recombinant protein production have been reported using COS, HEK 293 and BHK21 cells (Blasey et al., 1996, *Cytotechnology*, 18, 138-192; Jordan et al., 1998, *Cytotechnology*, 26, 39-47; Wurm and Bernard, 1999, *Curr Opin Biotechnol* 10, 156-159; Meissner et al., 2001, *Biotechn and Bioeng*, 75, 2, 197-203).

Transient expression can be performed by infection (using, *e.g.*, viral vectors) or by transfection of the appropriate vector. Transfection can be performed with either a non-replicating or a replicating DNA vector or with a virus derived RNA vector.

5 A variety of virus vectors for expression in higher eukaryotic cells have been developed by using recombinant DNA technology and introducing genes of interest into virus genomes. The initial viruses used in vector development were DNA viruses such as Adeno-associated viruses or Poxviruses as well as RNA viruses replicating through DNA intermediates such as Retroviruses.

10 Episomal replicating vectors provide certain advantages over classical expression systems. Some DNA viruses, such as Adeno-, Papilloma-, Polyoma-, Hepadna-, and Herpesviruses, do not normally integrate into the host genome, but replicate episomally (extrachromosomally) in the nucleus of a host cells such as in a mammalian host cell. This process may involve both virus trans-acting factors and
15 the host cell replication machinery. Episomal replicating vectors derived from these viruses generally contain a replication origin and at least one viral protein, *e.g.*, an initiator protein. Examples of such initiator proteins are large T-antigen for SV40, E1/E2 for BPV, and EBNA-1 for EBV.

Some host factors are able to recognize viral origins of replication and initiate
20 replication. For example human S/MAR (scaffold/matrix attached region from the human interferon-beta gene) is able to maintain the episomal replication of a vector containing SV 40 origin of replication. On the other hand certain viral origins of replication (ORI) are recognized by corresponding specific virally-encoded proteins. For example, SV40 or polyoma virus ORI is recognized by large T antigen, BPV
25 (Bovine papilloma virus) ORI is recognized by E1/E2 complex, EBV (Epstein-Barr virus) ORI is recognized by EBNA1 protein, and yeast origin of replication (ARS element) is recognized by ORE (origin recognition element) replication (Bode et al., 2001 Gene Ther Mol Biol, v6, 33-46).

The double strand DNA genome of Epstein-Barr virus EBV (165 kb) is
30 maintained in the latently infected cells as a large circular episome. Amplification of this episome is mediated by a cis-acting sequence such as the origin of replication (OriP), and one of the viral proteins, the Epstein-Barr viral nuclear antigen 1

(EBNA1). In a minimal system the presence of both OriP (1.8 kb region) and EBNA-1 is sufficient for replication and stabilization of recombinant plasmids. The exact function of EBNA-1 is still not clear, although the protein has a DNA binding domain. Generally EBNA-1 is involved in plasmid maintenance after completion of the synthesis and in plasmid segregation by dividing cells during proliferation. The OriP region is recognized also directly by the cellular DNA synthesizing machinery and is replicated during the cell cycle.

Other gene expression systems are based on alphaviruses (Lundstrom, K., *Curr. Opin. Biotechnol.* 8:578-582 (1997)). Several members of the alphavirus family, e.g., Sindbis virus (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989); Schlesinger, S., *Trends Biotechnol.* 11:18-22 (1993)), SFV (Liljeström, P. & Garoff, H., *Bio/Technology* 9:1356-1361 (1991)) and others (Davis, N.L. *et al.*, *Virology* 171:189-204 (1989)) have received considerable attention for the use as virus-based expression vectors for a variety of different proteins (Lundstrom, K., *Curr. Opin. Biotechnol.* 8:578-582 (1997); Liljeström, P., *Curr. Opin. Biotechnol.* 5:495-500 (1994)).

Alphaviruses are positive stranded RNA viruses which replicate their genomic RNA entirely in the cytoplasm of the infected cell and without a DNA intermediate (Strauss, J. and Strauss, E., *Microbiol. Rev.* 58:491-562 (1994)). The concept that alphaviruses can be developed as expression vectors was first established nearly ten years ago (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989)). Since then, several improvements have made the use of these RNA replicons as expression vectors more practical (Lundstrom, K., *Curr. Opin. Biotechnol.* 8:578-582 (1997)).

DNA vectors have been developed for both Sindbis virus (Herweijer, H. *et al.*, *Hum. Gene Ther.* 6:1495-1501 (1995); Dubensky, T.W. *et al.*, *J. Virol.* 70:508-519 (1996)) and SFV (Berglund, P. *et al.*, *Trends Biotechnol.* 14:130-134 (1996)). Eukaryotic promoters are introduced in these vectors upstream from the alphavirus replicase gene (consisting of the four non-structural protein genes (nsP1-4)), which are translated as one or two polyproteins which are then proteolytically cleaved. DNA is transcribed into RNA from the recombinant eukaryotic promoter in the nucleus and transported to the cytoplasm, where the replicase catalyzes the

replication of the alphavirus RNA molecule as during normal replication of the alphavirus RNA molecule (Strauss, J. and Strauss, E., *Microbiol. Rev.* 58:491-562 (1994)). Only transient expression of heterologous sequences has been possible until recently due to the cytopathogenicity of the alphavirus replicase (Lundstrom, K.,
5 *Curr. Opin. Biotechnol.* 8:578-582 (1997)).

Among the features which make Alphavirus vectors attractive as transient expression systems are the easy construction of plasmid clones, a rapid gene expression (within 24h), a high level of expression, a wide host range of the virus (many insect, avian and mammalian cells), production of high-titered stocks of
10 infectious particles and low risk of cell transformation due to the RNA nature of the genome and its replication entirely in the host cell cytoplasm without going through DNA intermediates. (Schlesinger S., 2000, *Exp. Opin. Biol. Ther.*, 1, 2, 177-191; Schlesinger S., *Adv. Virus res.*, v55, 565-577; Garoff, H. and K.J. Li, 1998, *Curr. Opin. Biotechnol.*, 9, 464-469; Lundstrom, K., 1997, *Curr. Opin. Biotechnol.*, 8, 578-
15 583; Xiong et al., 1989, *Science*, 243, 1188-1191). Blasey et al. compared the Semliki Forest Virus (SFV) expression system with the Epstein-Barr Virus (EBV) system (pCEP4 and HEK293EBNA cells) for expression of 5-HT3 receptors and found that both systems allow the production of functional recombinant protein within days to weeks at very high comparable levels (in: *Animal Cell Technology*,
20 1999, pp331-337).

One of the variants of alphavirus vectors is the replicon, which is capable of self-replication in the host cytoplasm after introduction into the cultured cells by transfection. In this replicon, viral structural protein genes have been replaced by a gene of interest (GOI). Such a replicon can be introduced into cells as RNA or cDNA
25 if placed under the control of an eukaryotic promoter or a promoter, which is recognized by the host cell machinery. (Agapov *et al.*, *Proc. Natl. Acad. Sci. USA* 95:12989-12994 (1998)).

RNA viruses have been engineered into chimeric viruses producing heterologous proteins. Such RNA virus vectors have been developed from
30 Alphaviruses (with positive strand genome), Influenza and VSV (with negative strand genome). One of the major advantages of these viruses is their RNA-only

replication strategy performed entirely in the cell cytoplasm and eliminating the risk of genome integration.

Adeno-, Adeno-associated and Retrovirus vectors can be used as gene therapy vectors and/or as expression vectors, especially for large-scale protein production. Herpesvirus derivative vectors containing a replication origin have been used for bioreactor protein production after transient transfection of a suitable cell line carrying a replication initiation gene. (Durocher *et al.*, *Nucleic Acids Res.* 30(2):E9 (2002); Meissner *et al.*, *Biotechnol. Bioeng.* 75(2):197-203 (2001)). In these experiments, 293E cells (293 human embryo kidney cells expressing Epstein-Barr virus nuclear antigen - EBNA-1) were used which provided a replication initiation factor in trans to a plasmid carrying EBV origin of replication – OriP. Reporter genes (encoding secreted, intracellular or transmembrane proteins) were constitutively expressed over a period of several days.

Hybrid vectors have also been reported, mainly as gene delivery systems or for use in virus biology research, such as combinations between EBV episomes and Herpes-, Adeno- (DNA) or Retro- (RNA-DNA) viruses (Sena-Esteves M. *et al.*, 1999, *Virol. Dec*;73(12):10426-39; Tan BT. *et al.*, 1999, *J Virol. Sep*;73(9):7582-9; Grignani F. *et al.*, 1998, *Cancer Res Jan 1*;58(1):14-9; Wang S. and Vos JM., 1996, *J Virol. Dec*;70(12):8422-30). EBV episomes are nuclear plasmids that are stably maintained through multiple cell divisions in primate and canine cells (J. L. Yates, N. Warren, and B. Sugden, 1985, *Nature* 313:812-815).

The ability to precisely control the expression of genes introduced into animal or human cells, or in whole organisms, enables significant progress in many areas of biology and medicine. For instance, methods that allow the intentional manipulation of gene expression will facilitate the analysis of genes whose expression cannot be tolerated constitutively.

To be of broad benefit, gene regulation techniques preferably allow for rapid, robust, precise and reversible induction of gene activity. As reviewed in Saez, E. *et al.*, (*Curr. Opin. Biotechnol.* 8:608-616 (1997)), an ideal system would fulfill the following requirements:

1. Specificity -- The system should be indifferent to endogenous factors and activated only by exogenous stimuli.

2. Non-interference -- The components of the system should not affect unintended cellular pathways.

3. Inducibility -- In the inactive state, the basal activity of the system should be minimal, while in the active state high levels of gene expression should be rapidly inducible.

4. Bioavailability of the inducer -- Inducing stimuli should rapidly penetrate to the site of interest.

5. Reversibility -- Inducing stimuli should clear swiftly to allow the system to rapidly return to the inactive state.

One common system currently in use for the regulation of gene expression is the tetracycline-based system (Gossen and Bujard, *Proc. Natl. Acad. Sci. USA* 89:5547 (1992)). This system is based on the continuous expression of a fusion protein with the tetracycline repressor protein (tetR) being converted into an activator by fusion to the transcriptional activation domain of the VP16 protein. The tTA system is useful for, *e.g.*, inducible gene expression and has been successfully used for the production of a number of proteins (Wimmel *et al.*, *Oncogene* 9:995 (1994); Früh *et al.*, *EMBO J.* 13:3236 (1994); Yu *et al.*, *J. Virol.* 70:4530 (1996)). However, there are certain disadvantages associated with the tTA system. (Schocket *et al.*, *Proc. Natl. Acad. Sci. USA* 92:6522 (1995); Howe *et al.*, *J. Biol. Chem.* 23:14168 (1995); Schocket and Schatz, *Proc. Natl. Acad. Sci. USA* 93:5173 (1996); Bohl *et al.*, *Nat. Med.* 3:299 (1997); Furth *et al.*, *Proc. Natl. Acad. Sci. USA* 91:9302 (1994); Hennighausen *et al.*, *J. Cell. Biochem.* 59:463 (1995); Kistner *et al.*, *Proc. Natl. Acad. Sci. USA* 93:10933 (1996); Hoffmann *et al.*, *Nucleic Acids Res.* 25:1078 (1997)).

Inducible expression systems based on viral expression vectors (*e.g.*, alphaviral vectors) are described in WO 99/50432 (incorporated herein by reference in its entirety). An exemplary alphaviral vector is pCytTs (also known as CytTs; *cf.* WO 99/50432). Using these systems, it is possible to precisely induce and regulate gene expression. This precise regulation results from the use of a temperature-sensitive RNA-dependent RNA polymerase (*i.e.*, a replicase), which only replicates RNA molecules, to form new RNA molecules, at permissive temperatures (Boorsma *et al.* *Nature Biotech.* 18:429-432 (2001)). Due to the extremely tight regulation of

these systems it is possible to obtain regulated gene expression even under conditions of transient transfection allowing the production of toxic proteins under these conditions.

Another example of an alphaviral expression vector is described in Lundstrom *et al.*, *FEBS Lett* 504:99-103 (2001). Lundstrom describe a rapidly generated high-titer Semliki Forest virus (SFV) vectors capable of infecting numerous mammalian cell lines and primary cell cultures, resulting in high levels of transgene expression. SFV-based expression of transmembrane receptors has been characterized by specific ligand-binding activity and functional responses. Adaptation of the SFV technology for mammalian suspension cultures has allowed the production of hundreds of milligrams of recombinant receptor for purification and structural studies. The same SFV stock solutions used for the infection of mammalian cells in culture have also been successfully applied for efficient transgene expression in organotypic hippocampal slices, as well as in vivo in rodent brain.

A need exists in the art, however, for compositions and methods that allow the production of polypeptides and untranslated RNA molecules.

BRIEF SUMMARY OF THE INVENTION

The present invention provides compositions and methods that allow the production of polypeptides and/or untranslated RNA molecules. More specifically, the invention provides nucleic acid molecules, expression systems, recombinant host cells that permit the production of polypeptides and/or untranslated RNA molecules. Also provided are methods for making the nucleic acid molecules, expression systems and host cells of the invention. The invention further provides methods for producing polypeptides and/or untranslated RNA molecules. The invention also provides kits that comprise the nucleic acid molecules, expression systems and/or recombinant host cells of the invention.

In certain embodiments, the invention provides compositions and methods for regulated, expression of polypeptides or untranslated RNA molecules in recombinant

host cells, preferably by way of transfection. The present invention also provides nucleic acid molecules and methods which allow, *e.g.*, rapid and high level production of specific RNA molecules produced in, *e.g.*, transiently transfected, recombinant host cells.

5 According to certain aspects of the invention, novel, preferably inducible, potent and rapid expression systems are provided comprising one or more nucleic acid molecules. The nucleic acid molecules of the expression systems of the invention may be maintained at, or accumulated to, multiple copies in the nuclei of recombinant host cells, preferably of transiently transfected mammalian cells. In
10 certain embodiments, RNA self replication may lead to the accumulation of a high number of RNA molecules in the cytoplasm of the transfected cells and subsequent translation of the polypeptide of interest. The nucleic acid molecules and expression systems of the invention are particularly useful for transient transfection.

 The present invention further provides nucleic acid molecules and methods
15 which allow inducible rapid and high level production of specific RNA molecules produced in, preferably transiently transfected, recombinant host cells. In certain embodiments, RNA self-replication is preferably inducible and triggered by a temperature shift. Thus, the high production levels, in certain embodiments, are achieved by the use of a temperature-sensitive RNA-dependent RNA polymerase
20 (*i.e.*, a replicase) which only replicates RNA molecules to form new RNA molecules at permissive temperatures. Moreover, if desired, the amount of RNA or polypeptide can be tightly regulated by the use of a temperature-sensitive RNA-dependent RNA polymerase. This facilitates, in particular, production of growth-inhibitory or toxic polypeptides or RNA-species.

25 In a general aspect, the present invention provides for a nucleic acid molecule comprising (A) a first polynucleotide element which encodes an RNA molecule, said RNA molecule comprising (a) at least one *cis*-acting sequence element; (b) a first nucleotide sequence comprising a first open reading frame, said first open reading frame having a nucleotide sequence encoding an RNA-dependent RNA polymerase;
30 and (c) at least one second nucleotide sequence selected from the group consisting of (i) a second open reading frame encoding a polypeptide; (ii) a nucleotide sequence complementary to all or part of the second open reading frame of (i); and (iii) a

nucleotide sequence encoding an untranslated RNA molecule or complement thereof; wherein said second nucleotide sequence is operably linked to a promoter which is recognized by said RNA-dependent RNA polymerase; (B) a second DNA element comprising an origin of replication; and (C) a third DNA element encoding a replication initiation factor capable of recognizing the origin of replication. Thus, the replication initiation factor preferably recognizes the origin of replication leading to replication of the second polynucleotide and second DNA element, respectively, comprising the origin of replication.

In certain embodiments, the RNA-dependent RNA polymerase is, *e.g.*, a non-cytopathic RNA-dependent RNA polymerase. In other embodiments, the RNA-dependent RNA polymerase is, *e.g.*, a temperature-sensitive RNA-dependent RNA polymerase. In yet other embodiments, the RNA-dependent RNA polymerase is, *e.g.*, a non-cytopathic, temperature-sensitive RNA-dependent RNA polymerase.

The nucleic acid molecules of the invention, in certain embodiments, comprise a 5' promoter which is capable of initiating transcription *in vivo*, 5' and/or 3' sequences enabling replication of the RNA molecule (*cis*-acting sequence elements), and a subgenomic promoter 5' to the gene of interest, as well as a sequence of interest which is translatable only after one or more RNA-dependent RNA replication events. These RNA-dependent RNA replication events are catalyzed by a, preferably regulatable, RNA-dependent RNA polymerase which may be encoded by the same mRNA molecule that is produced by transcription of the DNA vector or by a different mRNA molecule.

The inventive nucleic acid molecules, expression systems and vector systems of the present invention thus allow for large-scale transient transfection and very rapid production of polypeptides eliminating the need of isolating stably transformed high producing cell clones. The preferred embodiment and hereby the inducibility of the vector makes the invention especially suitable for the production of cytotoxic polypeptides, *e.g.*, polypeptides that are detrimental to the viability of a host cell.

In a preferred embodiment of the present invention, the origin of replication and the replication initiation factor are derived from a DNA virus. For example, the origin of replication may be derived from a herpes virus and the replication initiation factor may be the Epstein-Barr virus nuclear antigen 1 (EBNA1). However,

replication initiation factors may also derive from other viruses or be cellular factors, which are capable of recognizing or recognize viral origins of replication. As indicated, in a further embodiment, the RNA-dependent RNA-polymerase is temperature-sensitive and temperature-regulated, respectively. Preferably, the RNA-dependent RNA-polymerase is of viral, more preferably of alphaviral origin.

Thus, in a preferred embodiment, the present invention provides a hybrid vector and vector system combining the advantages of EBV episomal replication with the advantages of RNA virus replication, wherein the replication takes place entirely in the cytoplasm. This type of hybrid vector would be characterized by the maintenance of multiple copies of DNA due to polynucleotide replication, typically plasmid replication, followed by the accumulation of full-length transcripts containing the viral replicon. These viral full-length transcripts are then further exponentially amplified in the cytoplasm by the viral replicase leading to the accumulation of RNA virus replicons and protein synthesis. The system is extremely valuable for production of toxic polypeptides if the RNA virus replication is controllable or inducible.

In a specific embodiment, the invention includes the combination of a Herpesvirus mini-replicon unit, i.e. the cis-acting replication origin OriP (Origin of replication P) and the cis- or trans-acting gene product – EBNA-1 (Epstein-Barr virus nuclear antigen 1, a replication triggering factor) with the tightly regulated temperature inducible alphaviral expression system pCytTS, as disclosed in WO 99/504332. Due to the presence of OriP and EBNA-1 the introduced novel vector is maintained episomally (extrachromosomally) in the form of several DNA copies in the cell nuclei. These multiple DNA copies may be transcribed from a promoter, typically a CMV or RSV promoter, into CytTS RNA-replicons, but they remain inactive unless cells are being shifted to a certain temperature. Only after temperature induction replicon replication takes place, followed by RNA accumulation and translation in the cell cytoplasm and production of the polypeptide of interest.

In a further general aspect, the present invention provides a vector system comprising one or more nucleic acid molecules, wherein said one or more nucleic acid molecules comprise (A) a first polynucleotide which encodes an RNA molecule, said RNA molecule comprising (a) at least one *cis*-acting sequence element; (b) a

first nucleotide sequence comprising a first open reading frame having a nucleotide sequence encoding an RNA-dependent RNA polymerase; and (c) at least one second nucleotide sequence selected from the group consisting of (i) a second open reading frame encoding a polypeptide; (ii) a sequence complementary to all or part of the second open reading frame of (i); and (iii) a sequence encoding an untranslated RNA molecule, or complement thereof; wherein said second nucleotide sequence is operably linked to a promoter which is recognized by said non-cytopathic RNA-dependent RNA polymerase; (B) a second polynucleotide comprising an origin of replication; and (C) a third polynucleotide encoding a replication initiation factor capable of recognizing the origin of replication. Thus, the replication initiation factor recognizes the origin of replication leading to replication of the second polynucleotide and second DNA element, respectively, comprising the origin of replication.

The invention further provides single- and multiple-vector systems for producing a polypeptide or untranslated RNA molecule. In a single-vector system, the first, second and third polynucleotide are present on the same nucleic acid molecule. In a multiple-vector system, the first, second and third polynucleotide elements are present on one or more separate nucleic acid molecules. In such a multiple vector system, at least one, preferably some or more preferably all vectors contain an origin of replication. Therefore, in preferred embodiments of the invention, the replication initiation factor and the origin of replication recognizing the replication initiation factor may be on the same nucleic acid molecule and plasmid, respectively, as it is the RNA-replicase or on a separate nucleic acid molecule and plasmid, respectively.

When sequences encoding the first and second open reading frame are present either on the same nucleic acid molecule or in the same vector (i.e., in a single-vector system), a region will preferably be present 5' to the second open reading frame which inhibits translation of this open reading frame.

If a temperature sensitive RNA-dependent RNA polymerase (e.g., a temperature-sensitive replicase) is included within the compositions and methods of the invention, the temperature-sensitive replicase may be "cold" or "hot" sensitive and thus will only efficiently catalyze RNA-dependent RNA replication at

temperatures either above or below the restrictive temperature. In one preferred embodiment, the compositions and methods of the invention comprise, or involve the use of, an RNA-dependent RNA polymerase that has replicase activity at temperatures below 34°C which is at least five fold greater than the replicase activity exhibited at 34°C or above. Preferably, the replicase activity at temperatures below 34°C is at least ten fold greater than the replicase activity exhibited at 34°C or above. In certain embodiments, the replicase activity at temperatures below 34°C is at least one hundred fold greater than the replicase activity exhibited at 34°C or above. Temperature-sensitive RNA-dependent RNA polymerases included within the present invention may, in certain embodiments, exhibit no (or undetectable) replicase activity at 34°C or above.

In another aspect, the nucleic acid molecules and expression systems of the invention encode one or more cytokine, lymphokine, tumor necrosis factor, interferon, toxic protein, prodrug converting enzyme, or other polypeptide.

In yet another aspect, the nucleic acid molecules and expression systems of the invention encode an untranslated RNA molecule, such as an antisense RNA molecule, tRNA molecule, rRNA molecule, or ribozyme.

The invention also provides methods for making recombinant host cells comprising introducing nucleic acid molecules of the invention into host cells. Further provided are recombinant host cells produced by the introduction of nucleic acid molecules of the invention. In one embodiment, one, some or all of these recombinant host cells contain one or more nucleic acid molecules that comprise an RNA-dependent RNA polymerase and/or the replication initiation factor. The replication initiation factor, in certain embodiments, may be stably integrated into the genome of the one, some or all host cells.

The invention further provides isolated nucleic acid molecules comprising polynucleotides which comprise the nucleotide sequence of SEQ ID NO:21 (pCytTs-OriP) and SEQ ID NO:22 (pCytTs-OPE).

The present invention also provides methods for producing polypeptides and untranslated RNA molecules in recombinant host cells, said methods comprising introducing a nucleic acid molecule of the invention into a host cell to produce a

recombinant host cell, and culturing the recombinant host cell under conditions suitable for expression of said polypeptide or untranslated RNA molecule. The methods of the invention may further comprise recovering said polypeptide or untranslated RNA molecule.

5 Methods are also provided for the expression of heterologous polypeptides, preferably the regulated expression of heterologous polypeptides, including cytokines, lymphokines, tumor necrosis factors, interferons, toxic polypeptides, and prodrug converting enzymes.

10 Further provided are polypeptides and untranslated RNA molecules produced by the methods of the invention.

15 The invention also provides methods for regulating the expression of a polypeptide or an untranslated RNA molecule, said method comprising introducing nucleic acid molecules of the invention into a host cell to produce a recombinant host cell and growing said recombinant host cell under suitable culture conditions. In certain embodiments, nucleic acid molecules encoding a temperature-sensitive RNA-dependent RNA-polymerase are used. When a temperature-sensitive RNA-dependent RNA-polymerase is used, the methods according to this aspect of the invention may further comprise changing the temperature of the recombinant host cell culture from either (i) a permissive temperature to a restrictive temperature, or (ii) a restrictive
20 temperature to a permissive temperature.

25 In certain embodiments, the methods of the invention involve introducing the nucleic acid molecules of the invention into prokaryotic or eukaryotic host cells to produce a recombinant host cell, and then culturing said recombinant host cell under suitable culture conditions. The recombinant host cells may be cultured, *e.g.*, in a serum-free or protein-free medium.

 The present invention also provides pharmaceutical compositions comprising nucleic acid molecules of the invention and a pharmaceutically acceptable carrier.

30 Also included within the invention are kits comprising the nucleic acid molecules of the invention. Kits of the invention may additionally or alternatively comprise one or more expression systems of the invention and one or more recombinant host cells of the invention.

BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 is a schematic representation of vector pCytTs-EGFP-OPE described in Example 2. The EGFP expression cassette can be replaced by any gene of interest (e.g. SEAP, EPO, IFN). CMV: Cytomegalovirus promoter, puro: puromycin resistance marker; amp: ampicillin resistance marker; ColE1: bacterial origin of replication; SG prom: subgenomic promoter; ORIP, EBV origin of replication; EBNA1, EBV nuclear antigen 1; GFP: green fluorescence protein; nsp1-4: genes coding for the viral non-structural proteins containing the nsp 4 (Pro728Ser) mutation and the nsp2 (Gly153Glu) mutation that renders the replicase temperature sensitive and non-cytopathic.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to compositions and methods that are useful, e.g., in the production of polypeptides and/or untranslated RNA molecules in host cells. The compositions and methods of the invention are also useful, e.g., for the isolation and/or purification of polypeptides, polypeptides and untranslated RNA molecules that are produced in host cells.

According to one aspect of the invention, nucleic acid molecules are provided. The nucleic acid molecules of the invention comprise: (A) a first polynucleotide element which encodes an RNA molecule, said RNA molecule comprising: (a) at least one *cis*-acting sequence element; (b) a first nucleotide sequence comprising a first open reading frame, said first open reading frame having a nucleotide sequence encoding an RNA-dependent RNA polymerase; and (c) at least one second nucleotide sequence selected from the group consisting of: (i) a second open reading frame encoding a polypeptide; (ii) a nucleotide sequence complementary to all or a part of the second open reading frame of (i); and (iii) a nucleotide sequence encoding an untranslated RNA molecule or complement thereof; wherein said second nucleotide sequence is operably linked to a promoter which is recognized by said RNA-dependent RNA polymerase; (B) a second polynucleotide element comprising an origin of replication; and (C) a third polynucleotide element

encoding a replication initiation factor capable of recognizing said origin of replication.

According to another aspect of the invention, expression systems are provided. The expression systems of the invention comprise one or more nucleic acid molecules, wherein said one or more nucleic acid molecules comprise: (A) a first polynucleotide element which encodes an RNA molecule, said RNA molecule comprising: (a) at least one *cis*-acting sequence element; (b) a first nucleotide sequence comprising a first open reading frame, said first open reading frame having a nucleotide sequence encoding an RNA-dependent RNA polymerase; and (c) at least one second nucleotide sequence selected from the group consisting of: (i) a second open reading frame encoding a polypeptide; (ii) a nucleotide sequence complementary to all or a part of the open reading frame of (i); and (iii) a nucleotide sequence encoding an untranslated RNA molecule or complement thereof; wherein said second nucleotide sequence is operably linked to a promoter which is recognized by said RNA-dependent RNA polymerase; (B) a second polynucleotide element comprising an origin of replication; and (C) a third polynucleotide element encoding a replication initiation factor capable of recognizing said origin of replication.

In the expression systems of the invention, said first, second and third polynucleotide elements may each be on a separate nucleic acid molecule. Alternatively, said first, second and third polynucleotide elements may each be on a single nucleic acid molecule. In certain embodiments, said first and second polynucleotide elements will be on a single nucleic acid molecule while said third polynucleotide element is on a different nucleic acid molecule. In certain other embodiments, said first and third polynucleotide elements will be on a single nucleic acid molecule, while said second polynucleotide element is on a different nucleic acid molecule. In certain other embodiments, said second and third polynucleotide elements will be on a single nucleic acid molecule, while said first polynucleotide element is on a different nucleic acid molecule.

According to another aspect of the invention, recombinant host cells and *in vitro* cell cultures comprising recombinant host cells are provided. The recombinant host cells of the invention comprise: (A) a first polynucleotide element which

encodes an RNA molecule, said RNA molecule comprising: (a) at least one *cis*-acting sequence element; (b) a first nucleotide sequence comprising a first open reading frame, said first open reading frame having a nucleotide sequence encoding an RNA-dependent RNA polymerase; and (c) at least one second nucleotide sequence selected from the group consisting of: (i) a second open reading frame encoding a polypeptide; (ii) a nucleotide sequence complementary to all or a part of the second open reading frame of (i); and (iii) a nucleotide sequence encoding an untranslated RNA molecule or complement thereof; wherein said second nucleotide sequence is operably linked to a promoter which is recognized by said RNA-dependent RNA polymerase; (B) a second polynucleotide element comprising an origin of replication; and (C) a third polynucleotide element encoding a replication initiation factor capable of recognizing said origin of replication.

According to another aspect of the invention, methods are provided for producing, or regulating the expression of, a polypeptide or untranslated RNA molecule. The methods according to this aspect of the invention comprise: (a) introducing a nucleic acid molecule or an expression system of the invention into a host cell to produce a recombinant host cell; and (b) culturing said recombinant host cell. The methods of the invention may further comprise recovering said polypeptide or untranslated RNA molecule.

According to another aspect of the invention, kits are provided comprising the nucleic acid molecules, expression systems and/or host cells of the invention.

In certain embodiments, said second open reading frame that is included within the compositions and methods of the invention is in a translatable format after one or more RNA-dependent RNA replication events. The expression "a translatable format" is intended to mean an RNA from which a protein can be made.

The RNA-dependent RNA polymerase that is included in, or used with, the compositions and methods of the invention may, in certain embodiments, be selected from the group consisting of: (a) a temperature-sensitive RNA-dependent RNA polymerase, (b) a non-cytopathic RNA-dependent RNA polymerase, and (c) a temperature-sensitive, non-cytopathic RNA-dependent RNA polymerase.

When the RNA-dependent RNA polymerase that is included in, or used with, the compositions and methods of the invention is temperature sensitive, the

temperature-sensitive RNA-dependent RNA polymerase may have replicase activity at temperatures below 34°C which is at least five fold, at least ten fold, at least twenty fold, at least thirty fold, at least forty fold, at least fifty fold, at least one hundred fold, or at least one thousand fold greater than the replicase activity exhibited at 34°C or above. For example, the temperature-sensitive RNA-dependent RNA polymerase may have replicase activity at 34°C that is at least five fold greater than the replicase activity exhibited at 29°C. The temperature-sensitive RNA-dependent RNA polymerase may, in certain embodiments, exhibit no or undetectable replicase activity (using methods of detection known in the art) at 34°C or above.

The RNA-dependent RNA polymerase that is included in, or used with, the compositions and methods of the invention may, in certain embodiments, be of viral origin. For example, the RNA-dependent RNA polymerase may be of alphaviral origin. In specific embodiments of the invention, the RNA-dependent RNA polymerase is derived from a Sindbis virus, a Semliki Forest virus or an Aura virus.

The RNA-dependent RNA polymerase may be derived from a virus selected from the group consisting of Bebaru virus, Cabassou virus, Chikungunya virus, Eastern equine encephalomyelitis virus, Fort Morgan virus, Getah virus, Kyzylagach virus, Mayoaro virus, Middleburg virus, Mucambo virus, Ndumu virus, Pixuna virus, Tonate virus, Trinita virus, Una virus, Western equine encephalomyelitis virus, Whartaroa virus, Venezuelan equine encephalomyelitis virus (VEE), and Ross River virus.

The origin of replication that is included in, or used with, the nucleic acid molecules, the expression systems, compositions and methods of the invention may be derived from a prokaryotic organism, a eukaryotic organism (*e.g.*, a yeast, insect or mammal), or a virus. For example, the origin of replication may be derived from a DNA virus, *e.g.*, a DNA virus that allows for episomal replication. The origin of replication may be derived from a DNA virus selected from the group consisting of Herpesvirus, Epstein-Barr virus (EBV), Papillomavirus, Polyomavirus, Adenovirus, and Hepadnavirus. In a specific embodiment, the origin of replication is *oriP*, derived from EBV.

The replication initiation factor that is included in, or used with, the compositions and methods of the invention is, in certain embodiments, capable of

operating as a plasmid maintenance factor. As used herein, the term "plasmid maintenance factor" is intended to mean a factor, which supports the distribution of the episomes to the daughter cells upon cell division. In the case of EBNA-1, this is achieved by the binding of the protein to the episome and, at the same time, to the chromosome, thereby leading to proper distribution of the episomes to the daughter cells. The replication initiation factor may be derived from a prokaryotic organism, a eukaryotic organism (*e.g.*, a yeast, insect or mammal), or a virus. For example, the replication initiation factor may be derived from a DNA virus, *e.g.*, a DNA virus that allows for episomal replication. The replication initiation factor may be derived from a DNA virus selected from the group consisting of Herpesvirus, Epstein-Barr virus (EBV), Papillomavirus, Polyomavirus, Adenovirus, and Hepadnavirus. In a specific embodiment, the replication initiation factor is the EBNA-1 protein, derived from EBV.

The origin of replication and the replication initiation factor that are included in, or used with, the compositions and methods of the invention may, in certain embodiments, be derived from the same organism or the same virus. Alternatively, the origin of replication and the replication initiation factor may be derived from different organisms or viruses.

The nucleic acid molecules and expression systems of the invention may further comprise a fourth polynucleotide element, wherein said fourth polynucleotide element comprises a selection marker. Any selection marker known in the art may be used, including, *e.g.*, markers that confer resistance to antibiotics. An exemplary selection marker is one that confers resistance to puromycin. Alternatively, selection markers may be used that confer resistance to hygromycin, gpt, neomycin, zeocin, ouabain, blasticidin, or bleomycin, or markers such as DHFR, hisD, trpB, or glutamine synthetase.

The second open reading frame that is included in, or used with, the compositions and methods of the invention may encode, in certain embodiments, any polypeptide of interest. For example, the second open reading frame may encode, *e.g.*, a cytokine, a lymphokine, a tumor necrosis factor, an interferon (*e.g.*, beta-interferon, including human beta-interferon), a toxic polypeptide, or a prodrug.

The second open reading frame that is included in, or used with, the compositions and methods of the invention may encode, in certain embodiments, an untranslated RNA molecule. Exemplary untranslated RNA molecules include, *e.g.*, an antisense RNA molecule, a tRNA molecule, an rRNA molecule and a ribozyme. Other useful untranslated RNA molecules are known in the art.

Thus, in certain exemplary embodiments, the invention provides compositions and methods which allow for the production of high amounts of specific RNA molecules produced in host cells upon their transient transfection with said polynucleotides. This high production results, *e.g.*, from the use of a self-replicating episomal DNA plasmid combined with an RNA-dependent RNA polymerase, which amplifies the mRNA of interest.

Expression, using the compositions and methods of the invention, may be regulated precisely, if necessary. Such precise regulation can be achieved, *e.g.*, from the use of a temperature-sensitive RNA-dependent RNA polymerase which only replicates RNA molecules to form additional RNA molecules at permissive temperatures.

According to the present invention, polypeptides and/or untranslated RNA molecules may be produced in transient transfection experiments due to the use of nucleic acid molecules comprising an origin of replication and DNA encoding a protein (a factor or a DNA-dependent replicase) that can recognize the origin of replication, thereby resulting in replication of the plasmid.

The expression systems of the invention, in certain embodiments, comprise alphavirus DNA vectors that can be used to create transiently transfected cell lines. Said alphavirus DNA vectors may carry genes encoding a non-cytopathic replicase and viral non-structural proteins, thereby being able to produce high amounts of polypeptide. An induction of expression is achieved when the activity of the temperature-sensitive replicase is switched on by reducing the incubation temperature of the transfected cells from 37°C to a temperature lower than 34°C. Protein expression in the host at 37°C is, preferably, below the level of detection.

DEFINITIONS

The following definitions are provided to clarify the subject matter, which the inventors consider to be the present invention.

5 As used herein, a nucleic acid molecule is a sequence of contiguous nucleotides (riboNTPs, dNTPs or ddNTPs, or combinations thereof) of any length which may encode a full-length polypeptide or a fragment of any length thereof, or which may be non-coding. As used herein, the terms "nucleic acid molecule," and "polynucleotide," "polynucleotide construct," and "polynucleotide element" may be
10 used interchangeably.

As used herein, the term "alphavirus" refers to any of the RNA viruses included within the genus *Alphavirus*. Descriptions of the members of this genus are contained in Strauss and Strauss, *Microbiol. Rev.*, 58:491-562 (1994). Examples of alphaviruses are selected from the group comprising Aura virus, Bebaru virus,
15 Cabassou virus, Chikungunya virus, Easter equine encephalomyelitis virus, Fort morgan virus, Getah virus, Kyzylagach virus, Mayoaro virus, Middleburg virus, Mucambo virus, Ndumu virus, Pixuna virus, Tonate virus, Trinita virus, Una virus, Western equine encephalomyelitis virus, Whataroa virus, Sindbis virus (SIN), Semliki forest virus (SFV), Venezuelan equine encephalomyelitis virus (VEE), and
20 Ross River virus.

As used herein, the term "purified" used in reference to a molecule means that the concentration of the molecule being purified has been increased relative to molecules associated with it in its natural environment. Naturally associated molecules include polypeptides, nucleic acids, lipids and sugars but generally do not
25 include water, buffers, and reagents added to maintain the integrity or to facilitate the purification of the molecule. For example, even if mRNA is diluted with an aqueous solvent during oligo dT column chromatography, mRNA molecules are purified by this chromatography if naturally associated nucleic acids and other biological molecules do not bind to the column and are separated from the subject mRNA
30 molecules.

As used herein, the term "isolated" used in reference to a molecule means that the molecule has been removed from its native environment. For example, a

polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated." Further, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Isolated
5 RNA molecules include *in vivo* or *in vitro* RNA replication products of DNA and RNA molecules. Isolated nucleic acid molecules further include synthetically produced molecules. Additionally, vector molecules contained in recombinant host cells are also isolated. Thus, not all "isolated" molecules need to be "purified."

As used herein, the phrase "low or undetectable," when used in reference to
10 gene expression level, refers to a level of expression which is either significantly lower than that seen when the gene is maximally induced (*e.g.*, at least five fold lower) or is not readily detectable by the methods used in the following examples section.

As used herein, the phrase "individual" refers to multicellular organisms and
15 includes both plants and animals. Preferred multicellular organisms are animals, more preferred are vertebrates, even more preferred are mammals, and most preferred are humans.

As used herein, the term "pCytTs" means any temperature sensitive, non-cytopathic inducible alphaviral expression system containing a functional promoter to drive the transcription of mRNA from the nucleic acid molecule of the invention.
20 Examples for such expression systems or vectors may be, *e.g.*, the vector pCytTs (SEQ ID NO: 1) containing a RSV promoter or the vector pCytTs2.1 (SEQ ID NO: 2) containing a CMV promoter. When combinatorial constructs with pCytTs are mentioned in the Examples, they generally contain a CMV promoter, even if the
25 abbreviation of the combinatorial constructs does – for the sake of simplicity - only refer to pCytTs and not to pCytTs2.1.

As used herein, the phrase "*cis*-acting" sequence refers to nucleic acid sequences to which a replicase binds to catalyze the RNA-dependent replication of RNA molecules. These replication events result in the replication of the full-length
30 and partial RNA molecules and, thus, the alphavirus subgenomic promoter is also a "*cis*-acting" sequence. *Cis*-acting sequences may be located at or near the 5' end, 3' end, or both ends of a nucleic acid molecule, as well as internally.

As used herein, the phrase "RNA-dependent RNA polymerase" refers to a polymerase which catalyzes the production of an RNA molecule from another RNA molecule. This term is used herein synonymously with the term "replicase."

5 As used herein, the phrase "non-infective packaged RNA molecules" refers to packaged RNA molecules which will essentially undergo only one round of host cell infection and are not pathogenic. These molecules are thus "infective" but only for a single infectious entry into a host cell and are not capable of reproducing to form additional infectious particles.

10 As used herein, the term "transcription" refers to the production of RNA molecules from DNA templates catalyzed by RNA polymerases.

As used herein, the phrase "RNA-dependent RNA replication event" refers to processes which result in the formation of an RNA molecule using an RNA molecule as a template.

15 As used herein, the term "vector" refers to an a polynucleotide construct, typically a plasmid or a virus, used to transmit genetic material to a host cell. Preferably, the term "vector" as used herein refers to an agent such as a plasmid, and even more preferably to a circular plasmid, A vector as used herein may be composed of either DNA or RNA. Preferably, a vector as used herein is composed of DNA.

20 As used herein, the term "heterologous sequence" refers to a second nucleotide sequence present in a vector of the invention. The term "heterologous sequence" also refers to any amino acid or RNA sequence encoded by a heterologous DNA sequence contained in a vector of the invention. Heterologous nucleotide sequences can encode polypeptides or RNA molecules normally expressed in the cell
25 type in which they are present or molecules not normally expressed therein (*e.g.*, Sindbis structural proteins).

As used herein, the phrase "untranslated RNA" refers to an RNA sequence or molecule which does not contain an open reading frame or encodes an open reading frame, or portion thereof, but in a format in which an amino acid sequence will not
30 be produced (*e.g.*, no initiation codon is present). Examples of such molecules are tRNA molecules, rRNA molecules, and ribozymes. Antisense RNA may be

untranslated but, in some instances antisense sequences can be converted to a translatable sense strand from which a polypeptide is produced.

As used herein, the phrase "temperature-sensitive" refers to an enzyme which readily catalyzes a reaction at one temperature but catalyzes the same reaction slowly or not at all at another temperature. An example of a temperature-sensitive enzyme is the replicase protein encoded by the pCytTs vector, which has readily detectable replicase activity at temperatures below 34°C and has low or undetectable activity at 37°C.

As used herein, the phrase "permissive temperature" refers to temperatures at which an enzyme has relatively high levels of catalytic activity.

As used herein, the phrase "restrictive temperature" refers to temperatures at which an enzyme has low or undetectable levels of catalytic activity. Both "hot" and "cold" sensitive mutants are known and, thus, a restrictive temperature may be higher or lower than a permissive temperature.

As used herein, the term "recombinant host cell" refers to a host cell into which one or more nucleic acid molecules of the invention have been introduced.

As used herein, the term "replication initiation factor" refers to a protein or a DNA sequence encoding such a protein that is able to recognize an origin of replication leading to replication of plasmids containing this origin of replication.

As used herein, the term "origin of replication" refers to a DNA sequence that is recognized by a replication initiation factor or a DNA replicase leading to replication of a plasmid containing the origin of replication. The expression "recognized by a replication initiation factor" is intended to mean that a replication initiation factor can physically interact with all or a portion of an origin of replication sequence, thereby causing or stimulating molecular mechanisms that ultimately cause all or a portion of the DNA molecule comprising the origin of replication to be replicated.

As used herein the term "inducible" means that in an inactive state, the basal activity of a system should be minimal, while in the active state high levels of gene expression should be rapidly inducible.

When the terms "one," "a," or "an" are used in this disclosure, they mean "at least one" or "one or more," unless otherwise indicated.

NUCLEIC ACID MOLECULES AND EXPRESSION SYSTEMS

Certain preferred embodiments of the present invention are based on the vectors used in the previously described pCytTs expression system (WO 99/50432, the disclosure of which is hereby incorporated by reference in its entirety). These vectors are constitutively transcribed in host cells to produce mRNA molecules having two open reading frames. These open reading frames, which may or may not be produced from the same nucleic acid molecule, encode a temperature-sensitive replicase and a heterologous gene of interest. The first open reading frame is translated to produce an RNA-dependent RNA polymerase. The second open reading frame, encoding all or part of one or more polypeptides of interest, is not translated until after at least one RNA-dependent RNA replication event.

The nucleic acid molecule of the present invention comprise, *inter alia*, at least one second nucleotide sequence. The second nucleotide sequence is preferably operably linked a promoter which is recognized by an RNA-dependent RNA polymerase. The second nucleotide sequence, in certain embodiments, may comprise a second open reading frame, or other nucleotide sequence of interest, as well as other elements. For instance, the second nucleotide sequence may comprise a 5' promoter which is capable of initiating synthesis of RNA *in vivo*, 5' and/or 3' sequences enabling replication of the RNA molecule (5' and 3' *cis* acting sequence elements), as well as a sequence of interest which is translatable only after at least one replication event. Replication is catalyzed by an RNA-dependent RNA polymerase which is encoded alternatively on the same or on a different mRNA molecule. The sequence of interest may be encoded in sense, plus (+), orientation downstream of a viral RNA promoter. Translation of the coding sequence of the gene of interest is inhibited by a 5' sequence which, in the case of the single-vector system, will generally be the replicase sequence. In the multiple-vector system, a 5' sequence can inhibit translation by having one or more short open reading frames with associated stop codons which lead to the detachment of ribosomes. Similarly, any sequence which inhibits the traveling or binding of ribosomes to the sequence of interest can be used as a 5' sequence which inhibits translation (Voet and Voet, BIOCHEMISTRY, John Wiley & Sons, Inc. (1990)).

Another method for preventing translation of nucleotide sequences in most biological systems involves the insertion of the sequence in an antisense direction. This method of inhibiting translation is based on the principle that translation will generally only occur after the replication of this minus (-) strand RNA into a plus strand having an open reading frame in a sense orientation. The translated sense strand is formed by RNA replication and serves as a template for ribosomes and protein synthesis. As shown previously (WO 99/50432) production of amino acid sequences can occur even when the gene of interest is inserted into the DNA molecule in an orientation which will result in the formation of antisense RNA sequence 3' to the subgenomic promoter. Thus, the second open reading frame may also comprise a sequence complementary to all or part of the second open reading frame described above and expression of the encoded amino acid sequence will still occur. When the production of an untranslated antisense RNA sequence is desired, the RNA molecule can be designed so that it will not serve as a template for protein synthesis. For example, the RNA can be designed so that an initiation codon is not present.

The second nucleotide sequence may alternatively or additionally comprise a nucleotide sequence encoding an untranslated RNA molecule or complement thereof. For example, RNA molecules directly produced by transcription of a DNA sequence of the invention may encode RNA sequences which are neither translated nor present in an antisense orientation. Examples of such untranslated RNA molecules include tRNA molecules, rRNA molecules, and ribozymes. A considerable number of ribozyme sequences with defined catalytic activities are known in the art (*see, e.g.,* Brown, J., *Nucleic Acids Res.* 26:351-352 (1998); Xie, Y. *et al., Proc. Natl. Acad. Sci. USA* 94:13777-13781 (1997); Lavrovsky, Y *et al., Biochem. Mol. Med.* 62:11-22 (1997); Chapman, K. and Szostak, J., *Chem. Biol.* 2:325-333 (1995)). Further, ribozymes have been used to "knockout" the expression of a specific gene in eukaryotic cells as part of a ribozyme-mediated, message deletion strategy (Xie, Y. *et al., Proc. Natl. Acad. Sci. USA* 94:13777-13781 (1997)). Additionally, alphaviral replicons have been used to express a functional ribozyme in mammalian cells (Smith S. *et al., J. Virol.* 71:9713-9721 (1997)). The regulated expression of such

ribozymes, and other untranslated RNA molecules, is thus within the scope of the present invention.

The nucleic acid molecules and expression systems of the invention also comprise a second polynucleotide element comprising an origin of replication. Origins of replication that can be used with the present invention include, *e.g.*, those that are derived from prokaryotic organisms, eukaryotic organisms (*e.g.*, yeasts, insects, or mammals) and/or viruses. For example, an origin of replication that may be used with the present invention is one that is derived from a DNA virus such as, *e.g.*, a Herpesvirus, a Papillomavirus, an Adenovirus, or a Hepadnavirus. In certain embodiments, the origin of replication that is used with the invention is a DNA virus that allows for episomal replication. In a preferred embodiment, the origin of replication is derived from Epstein-Barr virus (EBV), such as, *e.g.*, *oriP*.

The nucleic acid molecules and expression systems of the invention also comprise a third polynucleotide element encoding a replication initiation factor. The replication initiation factor will be capable of recognizing the origin of replication that is included within the nucleic acid molecule or expression system. The third polynucleotide element of the invention may encode, *e.g.*, a replication initiation factor derived from prokaryotic organism, eukaryotic organism (*e.g.*, yeasts, insects, and/or mammals) and/or viruses. For example, a replication initiation factor that may be used with the present invention is one that is derived from a DNA virus such as, *e.g.*, a Herpesvirus, a Papillomavirus, an Adenovirus, or a Hepadnavirus. In certain embodiments, the replication initiation factor that is used with the invention is capable of operating as a plasmid maintenance factor. In a preferred embodiment, the replication initiation factor is derived from Epstein-Barr virus (EBV), such as, *e.g.*, the EBNA-1 protein, or a portion thereof.

The replication initiation factor (*e.g.*, a nucleotide sequence that encodes a replication initiation factor) and the origin of replication that are included within the nucleic acid molecules and expression systems of the present invention may be derived from the same organism or from the same virus. Alternatively, the replication initiation factor and the origin of replication may be derived from different organisms or from different viruses.

In an exemplary embodiment of the invention, nucleic acid molecules are provided, which are constructed by introducing defined Epstein-Barr virus (EBV) sequences into vectors such as, *e.g.*, those included within the pCytTs system. The resulting exemplary nucleic acid molecules of the invention are capable of replicating as non-integrated autonomous episomal molecules in the transformed host cells.

The defined sequences, which may be introduced into the vectors of the pCytTs system, comprise, *e.g.*, either one or two elements of Epstein-Barr virus, (OriP alone or together with EBNA-1 gene) which permit plasmid maintenance. The origin of replication OriP is a cis-acting sequence and needs to be inserted into the plasmid vector sequence. Plasmids containing this origin of replication are able to be maintained in cells expressing the replication initiation factor which recognizes the origin of replication; one of these factors is EBNA-1. EBNA-1 gene function can be provided in cis by introducing the sequence into the same plasmid or can be provided in trans by co-transfection with a second replicating or non-replicating plasmid as well as by providing it from stably transduced cell lines expressing EBNA-1 gene from an integrated copy. Furthermore, in certain embodiments, more than one copy of the sequence expressing the replication initiation factor, *e.g.* EBNA-1, is provided. For example, EBNA-1 gene function may be provided on one or more plasmids, wherein none, some or all of those plasmids may additionally comprise the origin of replication, and which plasmids are used for transfecting cell lines, preferably cell lines that have the sequence of EBNA-1 stably integrated within its genome. The presence of both OriP and EBNA1 sequences in the same plasmid renders replication less dependent on the host cell type. (Reisman, D. *et al.*, *Mol. Cell Biol.* 5: 1822-1832 (1985); Yates, J.L. *et al.*, *Nature* 313:812-815 (1985); U.S. Patent No. 4,686,186). U.S. Patent No. 4,686,186 describes the transfection of cells with a single plasmid containing the EBV OriP, the EBNA-1 gene and a gene encoding a protein of interest.

Although such cell lines have the advantage of stable long-term expression of the replication initiation factor and durable support of replication and maintenance of OriP containing plasmids, there are not many EBNA-1 expressing cell lines commercially available (ATCC : 293HEK-EBNA1 and CV1-EBNA1). Alternatively,

plasmids that already carry the EBNA-1 gene and the gene of interest in cis on the same episome are used to transfect cells and commercial vectors such as pCEP4 (Invitrogen) are available. However, current vectors designed for constitutive expression of polypeptides carrying OriP and EBNA-1 on the episomal construct in cis and wherein the expression is not regulatable or inducible, are not applicable in cases where cell-toxic polypeptides need to be expressed.

To our knowledge, an inducible expression system having the features of a self-replicating DNA episome and that allows the expression of a polypeptide or untranslated RNA sequence of interest from a self-replicating RNA molecule, triggered by a simple temperature shift, has not been described.

Certain embodiments of the invention are directed to nucleic acid molecules which are transcribed to produce a mRNA molecule having two open reading frames, *e.g.*, one open reading frame which encodes a replicase, and another open reading frame which encodes a nucleotide sequence of interest. The nucleic acid molecules of the invention may contain a promoter sequence which drives transcription to produce mRNA molecules having coding sequences of both open reading frames. The mRNA sequences of the first open reading frame are preferably translated to produce a replicase required for the expression of the RNA sequences of the second open reading frame. The second open reading frame preferably encodes one or more polypeptides of interest. In addition, the vectors contain an origin of replication and a DNA polymerase recognizing the origin of replication.

Further, once the first mRNA molecule has been transcribed from the DNA vector, additional RNA-dependent RNA replication events may occur to amplify the first mRNA sequence and to produce RNA molecules with strand polarity which is the opposite of the first mRNA sequence.

The second open reading frame of the nucleic acid molecules of the invention will preferably only be expressed after partial replication of a full-length RNA molecule. This partial replication of the full-length RNA molecules is driven by a promoter sequence composed of RNA (*e.g.*, an alphaviral subgenomic promoter sequence). In addition, the plasmid encoding the replicase and the gene of interest will preferably replicate during division of the host cell due to the presence of origin of replication and a protein factor recognizing the origin of replication.

While the gene of interest may be encoded by the same RNA molecule as the RNA-dependent RNA replicase protein, this gene may also be encoded by a separate RNA molecule. Thus, the invention further provides both single- and multiple-vectors systems for expressing a gene of interest. Moreover, while all plasmids used preferably contain an origin of replication, the gene encoding the replication initiation factor may be on a separate plasmid or stably integrated into the genome of the host cell.

In a single-vector system of the invention, sequences encoding the first open reading frame, the second nucleotide sequence are components of the same nucleic acid molecule, the origin of replication and the initiation factor. Thus, all of the components required for regulated expression of the gene of interest and plasmid amplification are contained in a single nucleic acid molecule (*i.e.*, DNA or RNA).

In a multiple-vector system of the invention, sequences encoding the first open reading frame, or sub-portions thereof, and the second nucleotide sequence are components of different nucleic acid molecules. These multiple-vector systems thus may comprise two or more nucleic acid molecules. For example, the replicase and the gene of interest can each be encoded by different nucleic acid molecules. However, all vectors may contain an origin of replication.

While any functional promoter can be used to drive the transcription of mRNA from the nucleic acid molecule of the invention, the promoter is preferably a constitutive RNA polymerase II promoter (*e.g.*, Rous Sarcoma Virus (RSV), cytomegalovirus (CMV), simian virus 40 (SV40), myeloproliferative sarcoma virus (MPSV), glucocorticoid, metallothionein, *Herpes simplex* virus thymidine kinase (HSVTK), human immune deficiency (HIV), mouse mammary tumor virus (MMTV), human polyomavirus BK (BKV), or Moloney murine leukemia virus (MuLV) promoter). Additional promoters suitable for use in the practice of the present invention are known in the art (*see, e.g.*, Lee, A. *et al.*, *Mol. Cells.* 7:495-501 (1997)).

The nucleic acid molecules of the invention may further comprise a fourth polynucleotide element, wherein said fourth polynucleotide element comprises a selection marker. The selection marker may facilitate the cloning and amplification of the vector sequences in prokaryotic and eukaryotic organisms. In certain

embodiments, the selection marker will confer resistance to a compound or class of compounds, such as an antibiotic. An exemplary selection marker that can be used with the nucleic acid molecules and expression systems of the present invention is one that confers resistance to puromycin. Alternatively, selection markers may be used that confer resistance to hygromycin, gpt, neomycin, zeocin, ouabain, blasticidin, or bleomycin, or markers such as DHFR, hisD, trpB, or glutamine synthetase.

The pCytTs vector contains an ampicillin resistance marker for positive selection in bacterial host cells and an *E. coli* origin of replication (*i.e.*, ColE1). A considerable number of sequences encoding additional selection markers and origins of replication are known in the art (*see, e.g.*, Sambrook, J. *et al.*, eds., MOLECULAR CLONING, A LABORATORY MANUAL, 2nd. edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Ausubel, F. *et al.*, eds., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John H. Wiley & Sons, Inc. (1997)).

The nucleic acid molecules and expression systems of the invention comprise a first nucleotide sequence comprising a first open reading frame, said first open reading frame having a nucleotide sequence encoding a replicase, *e.g.*, an RNA-dependent RNA polymerase. The first nucleotide sequence may, in certain embodiments, also comprise 5' and 3' *cis*-acting sequences, and junction sequences containing a subgenomic promoter.

The RNA-dependent RNA polymerases that can be used with the present invention may be of viral origin. The RNA-dependent RNA polymerase may be derived, *e.g.*, from virus selected from the group consisting of: Bebaru virus, Cabassou virus, Chikungunya virus, Easter equine encephalomyelitis virus, Fort morgan virus, Getah virus, Kyzylagach virus, Mayoaro virus, Middleburg virus, Mucambo virus, Ndumu virus, Pixuna virus, Tonate virus, Trinita virus, Una virus, Western equine encephalomyelitis virus, Whataroa virus, Venezuelan equine encephalomyelitis virus (VEE), and Ross River virus. In a preferred embodiment, the RNA-dependent RNA polymerase of the invention is derived from an alphavirus such as, *e.g.*, a Sindbis virus, a Semliki Forest virus, and/or an Aura virus.

In certain embodiments of the invention, the RNA-dependent RNA polymerase that is encoded by the first open reading frame of the nucleic acid

molecules of the invention will be a non-cytopathic RNA-dependent RNA polymerase. The RNA-dependent RNA polymerase may, alternatively, be a temperature sensitive RNA-dependent RNA polymerase. In other embodiments, the RNA-dependent RNA polymerase may be a non-cytopathic, temperature-sensitive
5 RNA-dependent RNA polymerase.

When the RNA-dependent RNA polymerase that is encoded by the first open reading frame of the nucleic acid molecules of the invention is a temperature-sensitive RNA-dependent RNA polymerase, the RNA-dependent RNA polymerase may, *e.g.*, have replicase activity at temperatures below 34°C and low or
10 undetectable replicase activity at 34°C or above. In certain embodiments, the temperature sensitive RNA-dependent RNA polymerase will be one that has replicase activity at 34°C which is at least five fold lower than the replicase activity exhibited at 29°C.

Preferred mutations which confer a non-cytopathic phenotype are those that
15 are in the nsp2 gene (*e.g.*, the proline residue at position 726 is replaced with a serine or a leucine residue). Mutations are known in the art which render the replicase protein non-cytopathic (Weiss *et al.*, *J. Virol.* 33:463-474 (1980); Dryga *et al.*, *Virology* 228:74-83 (1997); Agapov *et al.*, *PNAS* 95:12989-12994 (1998)). In addition, mutations may be introduced into the nuclear localization signal of an
20 alphaviral replicase which diminish toxicity of the replicon by, *e.g.*, modifying the distribution of the replicase-encoding sequence between the cytoplasm and the nucleus. (Rikkonen *et al.*, *Virology* 218:352-361 (1996)).

Mutations which render a replicase non-cytopathic may be introduced by a number of means, including site directed mutagenesis. Additional mutations that
25 render an RNA-dependent RNA polymerase non-cytopathic can be identified by, *e.g.*, random mutagenesis and screening using methods and techniques that are well-known in the art. As will be understood by persons having ordinary skill in the art, the identification of additional mutations that render an RNA-dependent RNA polymerase non-cytopathic, temperature sensitive, or both, does not require an ability
30 to predict the structural or functional consequences of any particular mutation.

About 20 years ago Weiss *et al.* (Weiss, B. *et al.*, *J. Virol.* 33:463-474 (1980)) established a persistently infected culture of BHK cells. The mutation

responsible for this phenotype has been recently identified (Dryga, S.A. *et al.*, *Virology* 228:74-83 (1997)). Another mutation allowing the regulation of the mRNA transcription via temperature shifts was identified by Burge and Pfefferkorn (Burge, B.W. & Pfefferkorn, E.R., *Virology* 30:203-214 (1966)) and described in more detail
5 by Xiong *et al.* (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989)).

Temperature sensitivity (*ts*) may be conferred, for example, by the introduction of a mutation in the *nsp4* gene of the replicase. Preferably, mutations which confer a temperature-sensitive phenotype upon replicase activities are in a protein in complementation group F (Lemm *et al.*, *J. Virol.* 64:3001-3011 (1990)).
10 For example, a temperature-sensitive phenotype may be conferred by changing Gly 153 of *nsp4* to Glu. Additionally, any other mutation which renders replicase activity temperature-sensitive can be used in the practice of the invention. Methods for creating and identifying new temperature-sensitive mutants are described by Pfefferkorn (Burge and Pfefferkorn, *Virol.* 30:204-213(1966); Burge and
15 Pfefferkorn, *Virol.* 30:214-223 (1966)). Other methods will be appreciated by those of ordinary skill in the art.

While most temperature-sensitive mutants are "hot" sensitive, "cold" sensitive mutations are also known (*see, e.g.*, Schwer, B. *et al.*, *Nucleic Acids Res.* 26:803-809 (1998), Mathe, E. *et al.*, *J. Cell Sci.* 111:887-896 (1998), Doedens, J. *et al.*, *J. Virol.* 71:9054-9064 (1997), Patterson, B. *et al.*, *J. Biol. Chem.* 272:27612-27617 (1997)). The temperature-sensitive replicase may be "cold" or "hot" sensitive and thus will catalyze RNA replication only at temperatures either above or below restrictive temperatures. In one embodiment, RNA replication occurs at detectable levels only at temperatures lower than 34°C. In contrast to all previously known
20 regulatable DNA expression systems, the basal level of expression in recombinant host cells containing the pCytTs vector in the inactive state at 37°C is below the level of detection using standard methods (*e.g.*, those used in the following examples) even under condition of transient transfection. This low level of expression is apparent from the data presented in Table 1.
25

30 The production of additional temperature sensitive and/or non-cytopathic replicases has been described in the art. (Lundstrom *et al.*, *Gene Ther. Mol. Biol.* 4:23-31 (1999), Lundstrom *et al.*, *Histochem. Cell. Biol.* 115:83-91 (2001)).

Moreover, with respect to alphaviruses, the results obtained in one type of alphavirus can often be used to predict the results obtained in other alphaviruses. That is, the identification of mutations that render a replicase temperature-sensitive and/or non-cytopathic in one alphavirus type may be used to identify or create a similar temperature-sensitive and/or non-cytopathic replicase in a different alphavirus. This ability is related, in part, to the high level of sequence identity observed among alphavirus genomes, especially in their replicase-encoding sequences. (Weaver *et al.*, *J. Virol* 71:613-623 (1997), Smyth *et al.*, *J. Virol.* 71:818-823 (1997), Kuhn *et al.*, *J. Virol.* 70:7900-7909 (1996)).

In certain embodiments, the first nucleotide sequence (comprising a first open reading frame encoding an RNA-dependent RNA polymerase) and the second nucleotide sequence (comprising, *e.g.*, a second open reading frame encoding a polypeptide) are contained on two separate nucleic acid molecules. In such an instance, the second nucleotide sequence may carry both *cis*-acting sequences and a 5' region which inhibits translation of the sequence of interest. The first nucleotide sequence can also be encoded by a nucleic acid molecule which is different than the second nucleotide sequence. Replication and translation of the second nucleotide sequence in this multi-vectors system can be regulatable by temperature if a temperature sensitive RNA-dependent RNA polymerase is used.

Additional mutations that render an RNA-dependent RNA polymerase non-cytopathic, temperature-sensitive, or both, can be identified by, *e.g.*, random mutagenesis and screening using methods and techniques that are well-known in the art. As will be understood by persons having ordinary skill in the art, the identification of additional mutations that render an RNA-dependent RNA polymerase non-cytopathic, temperature sensitive, or both, does not require an ability to predict the structural or functional consequences of any particular mutation. The identification of an RNA-dependent RNA polymerase that is non-cytopathic, temperature-sensitive, or both, only involves, *e.g.*, the generation of random mutations and the functional testing of the replicases that are encoded from the mutated nucleic acid molecules. Such screening methods are routine in the art.

The nucleic acid molecules and expression systems of the invention can be also used to express of more than one gene of interest. For example, recombinant

host cells can be transfected with more than one nucleic acid molecule of the invention wherein one nucleic acid molecule encodes both the replicase and a polypeptide of interest and additional nucleic acid molecules may encode additional polypeptides of interest. Similarly, when mutations conferring non-cytopathicity and temperature sensitivity are both used, genes encoding polypeptides having suitable mutations (*e.g.*, Pro 726 Ser in nsp2 and Gly 153 Glu in nsp4) may be on separate nucleic acid molecules. Additional variations would be apparent to those skilled in the art.

The nucleic acid molecules of the invention may also contain packaging signals which direct the packaging of RNA molecules into viral particles. These RNA molecules can be packaged in the presence of wild-type virus or defective helper virus RNA. A significant improvement was made with the development of defective helper RNA molecules (Bredenbeek, P. *et al.*, *J. Virol.* 67:6439-6446 (1993)). These RNA molecules contain *cis*-acting sequences, required for replication of the full-length transcription product, and subgenomic RNA promoter sequences which drive the expression of the structural protein genes. For example, in cells containing both RNA molecules with packaging signals and the defective helper virus RNA, alphaviral non-structural proteins allow for replication and amplification of the defective helper virus RNA sequences which are translated to produce virion structural proteins. Since the helper virus RNA lacks packaging signals, these molecules are not packaged into assembled virions. Thus, virion particles produced in this way contain essentially only RNA sequences encoding the gene of interest and, generally, other sequences required for temperature-sensitive regulation of gene expression. These non-infective packaged RNA molecules do not contain sequences encoding virion structural proteins and, thus, undergo only one round of host cell infection and are not pathogenic.

Non-infective packaged RNA molecules can be used to infect a culture of suitable host cells simply by adding the particles to culture medium containing these cells. The preparation of non-infective alphaviral particles is described in a number of sources, including "Sindbis Expression System", Version C, (Invitrogen Catalog No. K750-1).

One exemplary application of this system is directed to the temperature-dependent production of non-infective, packaged RNA molecules. These packaged RNA molecules may be produced by a number of means including using recombinant host cells containing two different nucleic acid molecules (*e.g.*, a nucleic acid molecule of the invention and a nucleic acid molecule encoding a helper virus RNA sequence). For example, one of these nucleic acid molecules will encode, *e.g.*, an RNA molecule which contains packaging signal sequences, sequences encoding a non-cytopathic, temperature-sensitive replicase, and the gene of interest. The other nucleic acid molecule will contain sequences encoding alphaviral structural proteins downstream from an alphavirus subgenomic promoter. Using such a system, viral particles containing only RNA molecules with packaging signals will be produced at permissive temperatures in recombinant host cells. This is so because alphaviral structural proteins will only be produced at a permissive temperature. Additional variations of the above would be apparent to one skilled in the art.

The second nucleotide sequence that is included within the nucleic acid molecules and expression systems of the invention may be, *e.g.*, a second open reading frame encoding a polypeptide. Such a second open reading frame may, alternatively or interchangeably, be referred to as a nucleotide sequence of interest. A wide variety of nucleotide sequences of interest can be expressed by the nucleic acid molecules and expression systems of the invention. These sequences include, but are not limited to, sequences encoding lymphokines, cytokines, toxins, enzymes, prodrug converting enzymes, antigens which stimulate immune responses, single chain antibodies, polypeptides which stimulate or inhibit immune responses, tumor necrosis factors, and various proteins with therapeutic applications (*e.g.*, growth hormones and regulatory factors).

In certain embodiments, the second open reading frame (the second nucleotide sequence) of the invention may encode, *e.g.*, a cytokine or lymphokine (*e.g.*, β -interferon). Hematopoiesis is regulated by lymphokines and cytokines which stimulate the proliferation and/or differentiation of various hemopoietic cells. Representative examples of cytokines and lymphokines include interleukin-1 (IL-1), interleukin-2 (IL-2), interleukin-3 (IL-3), interleukin-4 (IL-4), interleukin-5 (IL-5),

interleukin-6 (IL-6), interleukin-7 (IL-7), interleukin-8 (IL-8), interleukin-9 (IL-9), interleukin-10 (IL-10), interleukin-11 (IL-11), interleukin-12 (IL-12), interleukin-13 (IL-13), interleukin-14 (IL-14), interleukin-15 (IL-15), interleukin-16 (IL-16), interleukin-17 (IL-17), granulocyte colony stimulating factor (G-CSF), granulocyte-
5 macrophage colony stimulating factor (GM-CSF), macrophage colony stimulating factor (M-CSF), and interferons.

In certain embodiments, the second open reading frame (the second nucleotide sequence) of the invention may encode secreted enzymes (*e.g.*, secreted alkaline phosphatase), cytoplasmic enzymes (*e.g.*, green fluorescent protein), or any
10 number of other proteins with therapeutic applications (*e.g.*, human insulin, human coagulation Factor VIII).

The nucleic acid molecules and expression systems of the invention can also be used to express heterologous sequence encoding cytotoxic polypeptides. Cytotoxic polypeptides act to directly or indirectly inhibit cellular growth or metabolism. Representative examples of toxins include *Shigella* toxin, ricin,
15 *Diphtheria* toxin, *Cholera* toxin, *Pseudomonas* exotoxin A, and *Herpes simplex* virus thymidine kinase (HSVTK). Within other embodiments of this invention, the heterologous sequence encodes a prodrug converting enzyme. A prodrug converting enzyme activates a compound with little or no cytotoxicity into a toxic product.
20 Representative examples are HSVTK, alkaline phosphatase, guanine phosphoribosyl transferase, and penicillin-V amidase. Examples of both cytotoxic polypeptides and prodrug converting enzymes are discussed in numerous sources including PCT/US97/06010, EP 0716148, and WO 96/17072. In addition, a vast array of signaling molecules and membrane proteins are toxic if expressed at high levels. All
25 these molecules may be suitable for expression using the system of the present invention.

The second nucleotide sequence of the nucleic acid molecules and expression systems of the invention may also be a nucleotide sequence encoding an untranslated RNA molecule or complement thereof. Exemplary untranslated RNA molecules
30 include, *e.g.*, antisense sequences, RNase P targeted sequences which induce gene down-regulation, and ribozymes. Smith S. *et al.* (*J. Virol.* 71:9713-9721 (1997)) describes alphaviral vectors used to express ribozyme sequences.

The nucleic acid molecules and expression systems of the invention can also be used to express virtually any polypeptide, including ones which have not as yet been identified but are encoded by nucleotide sequences contained in, for example, cDNA libraries or host cell chromosomes. Example of such polypeptides include secreted proteins and proteins from various cellular compartments. Heterologous sequences expressed by the vectors of the invention can encode polypeptides and RNA molecules from non-human species (*e.g.*, other mammals, plants, fungi, bacteria or viruses). These heterologous sequences may further encode viral membrane proteins (*e.g.*, HIV gp160) or viral polyproteins (*e.g.*, Sindbis structural proteins).

Nucleotide sequences may be added to the nucleic acid molecules and vectors of the invention which result in the production of a fusion protein. For example, such sequences can encode amino acids sequences which are fused to a protein encoded by a gene of interest and confer one or more functional characteristics upon the expression product. These amino acid sequences include sequences which will target the gene product for export from the cell (*e.g.*, a secretory sequence) or to a subcellular compartment (*e.g.*, the nucleus). Such amino acid sequences further include sequences which facilitate purification (*e.g.*, a six His "tag"). Depending on the amino acid sequence and the function imparted by the fused sequence, the added amino acid sequences may or may not be cleaved from the translation product.

Fusion proteins also include proteins which have domains or regions derived from various different proteins. Examples of such a fusion protein are those containing domain II of *Pseudomonas* exotoxin, a domain or amino acid sequence which has binding affinity for a cell surface receptor associated with a particular cell type, and another amino acid sequence with a preselected biological activity. Domain II of *Pseudomonas* exotoxin will translocate across cell membranes. Using this system, fusion proteins can be designed which will bind to specific cells types, will translocate across the cytoplasmic membranes of these cells, and will catalyze predetermined intracellular biological reactions. A system of this type is described in Pastan *et al.*, U.S. Patent No. 5,705,163. Methods for identifying amino acid sequences, which bind to specific cell types are described in Wu, A., *Nature Biotech.* 14:429-431 (1996).

The nucleic acid molecules and expression systems of the invention may, in certain embodiments, further comprise genetic elements which confer additional functional characteristics such as selection markers.

5 Markers for the selection of prokaryotic and eukaryotic cells containing vectors the present invention are well known in the art and include puromycin, tetracycline, ampicillin, neomycin, hygromycin, gpt, zeocin, ouabain, blasticidin, bleomycin, and kanamycin resistance. Alternatively, selection markers such as DHRF, hisD, trpB, or glutamine synthetase may be used. Nucleotide sequences which result in high copy number amplification are also known in the art and include
10 the ColE1 sequence contained in the pCytTs vector.

RECOMBINANT HOST CELLS

15 The invention includes methods of making recombinant host cells and recombinant host cells produced using the methods of the invention. For example, the methods of the invention comprise introducing one or more nucleic acid molecules or expression systems described herein into a host cell. A variety of different recombinant host cells can be produced which contain the nucleic acid
20 molecules and expression systems of the invention. Alphaviruses, for example, are known to have a wide host range. Sindbis virus, for example, infects cultured mammalian, reptilian, and amphibian cells, as well as some insect cells (Clark, H., *J. Natl. Cancer Inst.* 51:645 (1973); Leake, C., *J. Gen. Virol.* 35:335 (1977); Stollar, V. in *THE TOGAVIRUSES*, R.W. Schlesinger, Ed., Academic Press, (1980), pp.583-621).

25 Thus, numerous host cells can be used in the practice of the invention. Representative host cells that may be used with the invention include, but are not limited to, bacterial cells, yeast cells, plant cells and animal cells. Preferred bacterial host cells include *Escherichia* spp. cells (particularly *E. coli* cells and most particularly *E. coli* strains DH10B, Stbl2, DH5, DB3, DB3.1, DB4 and DB5),
30 *Bacillus* spp. cells (particularly *B. subtilis* and *B. megaterium* cells), *Streptomyces* spp. cells, *Erwinia* spp. cells, *Klebsiella* spp. cells, *Serratia* spp. cells (particularly *S. marcescans* cells), *Pseudomonas* spp. cells (particularly *P. aeruginosa* cells), and

Salmonella spp. cells (particularly *S. typhimurium* and *S. typhi* cells). Preferred animal host cells include insect cells (most particularly *Drosophila melanogaster* cells, *Spodoptera frugiperda* Sf9 and Sf21 cells and *Trichoplusia* High-Five cells), nematode cells (particularly *C. elegans* cells), avian cells, amphibian cells (particularly *Xenopus laevis* cells), reptilian cells, and mammalian cells (most particularly NIH3T3, CHO, COS, VERO, BHK, HEK, other rodent cells, and human cells). Preferred yeast host cells include *Saccharomyces cerevisiae* cells and *Pichia pastoris* cells. BHK, COS, Vero, HeLa and CHO cells are particularly suitable for the production of heterologous polypeptides because they have the potential to glycosylate heterologous proteins in a manner similar to human cells (Watson, E. *et al.*, *Glycobiology* 4:227, (1994)) and can be selected (Zang, M. *et al.*, *Bio/Technology* 13:389 (1995)) or genetically engineered (Renner W. *et al.*, *Biotech. Bioeng.* 47:476 (1995); Lee K. *et al.* *Biotech. Bioeng.* 50:336 (1996)) to grow in serum-free medium, as well as in suspension.

Example 2 shows that the expression system of the present invention can function in BHK cells and 293HEK cells. Moreover, we have previously demonstrated that the pCytTs system works efficiently in additional cells lines such as CHO-K1 and COS-7 (Boorsma *et al.*, *Nature Biotech.* 18:429-432 (2000)).

The nucleic acid molecules and/or expression systems of the invention may be introduced into host cells using well known techniques of infection, transduction, electroporation, transfection, and transformation. Exemplary methods include DEAE-dextran mediated transfection, transient transfection, microinjection, cationic lipid-mediated transfection, scrape loading and ballistic introduction. Methods for the introduction of exogenous DNA sequences into host cells are discussed in Felgner, P. *et al.*, U.S. Patent No. 5,580,859. The nucleic acid molecules and/or vectors of the invention may be introduced alone or in conjunction with other nucleic acid molecules and/or vectors and/or proteins, peptides or RNAs. Alternatively, the nucleic acid molecules and/or expression systems of the invention may be introduced into host cells as a precipitate, such as a calcium phosphate precipitate, or in a complex with a lipid. Electroporation also may be used to introduce the nucleic acid molecules and/or expression systems of the invention into a host. Likewise, such molecules may be introduced into chemically competent cells such as *E. coli*. If the

vector is a virus, it may be packaged in vitro or introduced into a packaging cell and the packaged virus may be transduced into cells. Hence, a wide variety of techniques suitable for introducing the nucleic acid molecules and/or vectors of the invention into host cells are well known and routine to those of skill in the art. Such techniques are reviewed at length, for example, in Sambrook, J., et al., Molecular Cloning, a Laboratory Manual, 2nd Ed., Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, pp. 16.30-16.55 (1989), Ausubel, F. *et al.*, eds., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John H. Wiley & Sons, Inc. (1997), Chapter 16), Watson, J.D., *et al.*, Recombinant DNA, 2nd Ed., New York: W.H. Freeman and Co., pp. 213-234 (1992), and Winnacker, E.-L., From Genes to Clones, New York: VCH Publishers (1987), which are illustrative of the many laboratory manuals that detail these techniques and which are incorporated by reference herein in their entireties for their relevant disclosures.

The invention is also directed to recombinant host cells comprising one or more nucleic acid molecules or expression systems of the invention. Also included within the present invention are *in vitro* cell cultures that comprise such recombinant host cells. Methods for producing recombinant host cells and cell cultures comprising the same are well-known in the art.

PRODUCTION OF POLYPEPTIDES AND RNA MOLECULES

The nucleic acid molecules, expression systems and recombinant host cells of the invention may be used for the production of proteins, polypeptides and RNA molecules, *e.g.*, untranslated RNA molecules. The methods of the invention may comprise, *e.g.*, introducing one or more nucleic acid molecules or expression systems of the present invention into host cells to produce recombinant host cells, culturing the recombinant host cells under conditions suitable for expression of the polypeptide or untranslated RNA molecule, and recovering the polypeptide or untranslated RNA molecule.

The invention also provides methods for regulating the expression of a polypeptide or an untranslated RNA molecule. The methods of the invention may

comprise, *e.g.*, (a) introducing one or more nucleic acid molecules or expression systems of the invention into a host cell to produce a recombinant host cell, wherein said nucleic acid molecules or expression systems comprise a first open reading frame having a nucleotide sequence encoding a temperature-sensitive RNA-
5 dependent RNA polymerase, (b) growing said recombinant host cell under suitable culture conditions, and (c) changing the temperature of the recombinant host cell culture from: (i) a permissive temperature to a restrictive temperature, or (ii) a restrictive temperature to a permissive temperature.

The present invention also provides methods for producing polypeptides and
10 RNA molecules, said methods comprising introducing one or more nucleic acid molecules or expression systems of the invention into recombinant host cells, expanding the cells at a non-permissive temperature and incubating these cells at a permissive temperature. In a related aspect, the invention provides purified polypeptides and RNA molecules produced according to the methods of the present
15 invention.

Depending on the molecule, which is expressed, it may be obtained either from the culture supernatant or by lysing the recombinant host cells.

Polypeptides produced using the nucleic acid molecules and expression systems of the invention can be recovered and purified from recombinant cell
20 cultures by methods known in the art including ammonium sulfate precipitation, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and high performance liquid chromatography. Methods for purifying proteins are described in numerous sources (*see, e.g.*, Celis, J., ed., CELL
25 BIOLOGY, Academic Press, 2nd edition, (1998)).

Untranslated RNA molecules produced using the nucleic acid molecules and expression systems of the invention can be recovered and purified from recombinant cell cultures by methods known in the art (*see, e.g.*, Celis, J., ed., CELL BIOLOGY, Academic Press, 2nd edition, (1998)). Methods for recovering and/or purifying RNA
30 molecules include phenol/chloroform extraction, digestion with DNAses followed by precipitation of the undigested RNA molecules, and column chromatography (*e.g.*, oligo dT column chromatography). Further, RNA molecules can be separated from

other cellular material using the single-step guanidinium-thiocyanate-phenol-chloroform method described in Chomczynski and Sacchi, *Anal. Biochem.* 162:156-159 (1987).

5 The overall cell culture process employing nucleic acid molecules and expression systems of the invention for the production of polypeptides and/or untranslated RNA molecules can be implemented in a variety of bioreactor configurations (*e.g.*, stirred-tank, perfused, membrane enclosed, encapsulated cell, fluidized bed, and air-lift reactors) and scales (from laboratory T-flasks to thousands of liters), chosen to accommodate the requirements of the host cell line utilized (*e.g.*,
10 anchorage dependency, O₂ concentrations), to maximize the production of expression product, and to facilitate subsequent recovery and purification of expression product.

The invention is also directed to the production of polypeptides or RNA molecules of interest using mammalian cells grown in serum-free or protein-free culture media. For example, by long-term culture under conditions restricting serum
15 access or selecting for suspension growth, CHO cell lines are selected which are able to grow in serum-free medium and/or in suspension (Zang. M. *et al.*, *Bio/Technology* 13:389 (1995)).

Further, a number of different bioprocess parameters can be varied in order to alter the glycosylation pattern of polypeptide products produced by the recombinant
20 host cells of the invention. These factors include medium composition, pH, oxygen concentration, lack or presence of agitation, and, for the case of anchorage-dependent cells, the surface provided. Thus, the glycosylation pattern of glycoproteins may be altered by choosing the host cell in which these proteins are expressed in and the conditions under which the recombinant host cells are grown.

25 PHARMACEUTICAL COMPOSITIONS

The invention further provides pharmaceutical compositions comprising
30 nucleic acid molecules and/or expression systems and/or recombinant host cells of the invention. The pharmaceutical compositions of the invention may comprise nucleic acid molecules and/or expression systems and/or recombinant host cells of

the invention in combination (*e.g.*, in solution) with a physiologically acceptable carrier and in a therapeutically effective amount. The administration of these pharmaceutical compositions may, for example, result in expression of a polypeptide in tissues of an animal which is immunogenic and intended to function as a vaccination. Similarly, the nucleic acid molecules and/or expression systems and/or recombinant host cells of the invention may carry sequences that encode polypeptides or RNA molecules required for the treatment of an active affliction. The administration of a pharmaceutical composition of the invention will thus be intended to have a therapeutic effect in these instances.

The nucleic acid molecules and/or expression systems and/or recombinant host cells of the invention will normally be administered to an individual in a pharmacologically acceptable carrier. A composition is said to be "pharmacologically acceptable" if its administration can be tolerated by a recipient individual. Further, the composition of the invention will be administered in a "therapeutically effective amount" (*e.g.*, an amount that produces a desired physiological effect).

As would be understood by one of ordinary skill in the art, when the nucleic acid molecules and/or expression systems and/or recombinant host cells of the invention are administered to an individual, they may be in a composition which contains salts, buffers, adjuvants, or other substances which are desirable for improving the efficacy of the composition. Examples of materials suitable for use in preparing pharmaceutical compositions are provided in numerous sources including REMINGTON'S PHARMACEUTICAL SCIENCES (Osol, A, ed., Mack Publishing Co., (1980)).

The pharmaceutical compositions of the present invention can be administered by various art known means but will normally be administered by injection, infusion or other suitable physical methods. The compositions may alternatively be administered intramuscularly, intravenously, or subcutaneously. Components of compositions for administration include sterile aqueous (*e.g.*, physiological saline) or non-aqueous solutions and suspensions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as

olive oil, and injectable organic esters such as ethyl oleate. Carriers or occlusive dressings can be used to increase skin permeability and enhance antigen absorption.

When recombinant host cells are administered to an individual, the number of cells or nucleic acid molecules required to provide a therapeutically effective amount will vary with such factors as the individual's condition, the polypeptides or RNA molecules intended to be expressed, and the size of the individual.

KITS

10

The invention also provides kits comprising the isolated nucleic acid molecules, expression systems and/or recombinant host cells of the invention. The kits of the invention may optionally comprise one or more additional components selected from the group consisting of one or more containers (*e.g.*, boxes, vials, tubes, jars ampules, etc.) one or more vectors, one or more nucleotides, one or more primers, one or more polypeptides having polymerase activity, one or more host cells (*e.g.*, host cells that may be competent for uptake of nucleic acid molecules), and one or more buffers.

It will be readily apparent to one of ordinary skill in the relevant arts that other suitable modifications and adaptations to the methods and applications described herein are obvious and may be made without departing from the scope of the invention or any embodiment thereof. Having now described the present invention in detail, the same will be more clearly understood by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting of the invention.

EXAMPLE 1

CONSTRUCTION OF THE pCytTs VECTOR SYSTEM

Manipulations and sequencing of DNA were carried out by standard procedures. The mutations in nsP2 were introduced by PCR using the following oligonucleotides:

oligo-nsp2 1: 5'-AACATTGAAATCGATATTACAGGGG (SEQ ID NO:3),

oligo-nsp2 2: 5'-CGGGTTATGGTCGACCGGGC (SEQ ID NO:4),

oligo-nsp2 3: 5'-GTGCCCTCCCCTGAGTTTAACAATTCAGGGCCGAA
CGCG (SEQ ID NO:5), and

oligo-nsp2 4: 5'-GAATTGTTTAAACTCAGGAGGCACCCTCGTGG (SEQ ID NO:6).

The single restriction sites used for first analysis and subsequent cloning (*Dra*I, *Cla*I and *Sal*I) are underlined. PCR reactions were performed using either oligo-nsp2 1 (SEQ ID NO:3) and oligo-nsp2 3 (SEQ ID NO:5) or oligo-nsp2 2 (SEQ ID NO:4) and oligo-nsp2 4 (SEQ ID NO:6). 100 pmol of each oligo was used and 5 ng of the template DNA (pSinRep5; Xiong, C. *et al.*, *Science* 243:1188-1191 (1989)) was used in the 100 Φ l reaction mixture, containing 4 units of Taq or Pwo polymerase, 0.1 mM dNTPs and 1.5 mM MgSO₄. All DNA concentrations were determined photometrically using the GeneQuant apparatus (Pharmacia Biotech Inc., 800 Centennial Ave., Piscataway, NJ. 08854). The polymerase was added directly before starting the PCR reaction (starting point was 95°C). The temperature cycles were as follows: 95°C for 2 minutes, followed by 5 cycles of 95°C (45 seconds), 58°C (30 seconds), 72°C (90 seconds) and followed by 25 cycles of 95°C (45 seconds), 68°C (30 seconds), 72°C (90 seconds).

The two PCR fragment were purified using Qia spin PCR kit (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311) and finally digested in an appropriate buffer using 20 units of *Sal*I and *Dra*I, respectively 20 units of *Cla*I and *Dra*I. The digestion was performed for 12 hours at 37°C. The DNA fragments were gel-

purified (Gene-Clean; Bio 101 Inc., 1070 Joshua Way, Vista, CA, 92083, USA) and finally ligated into *ClaI/SalI* digested and gel-purified SinRep5 vector (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989). The correct sequence of the obtained vector was checked by DNA sequencing of the whole nsP2 gene.

5 The mutations in nsP4 were also introduced by PCR using the following oligonucleotides:

 oligo-nsP4 1: 5'-GGTAGACGAGACAGTCGCATGCCCTGGATAC (SEQ ID NO:7),

 oligo-nsP4 2: 5'-GTATCCAGGCGATGCGACTGTCTCGTCTACC (SEQ ID NO:8),
10

 oligo-nsP4 3: 5'-CAGACCGGTTAACGCCATAGCG TCG (SEQ ID NO:9),
and

 oligo-nsP4 4: 5'-CTCTATTACTAGTATGGACAGTTGG (SEQ ID NO:10),

 The singular restriction sites used for the first analysis and the final cloning
15 step (*SphI*, *HpaI* and *SpeI*) are underlined. Two PCR reactions were carried out as described above using either oligo-nsP4 1 (SEQ ID NO:7) and oligo-nsP4 3 (SEQ ID NO:9) or oligo-nsP4 2 (SEQ ID NO:8) and oligo-nsP4 4 (SEQ ID NO:10).

 Both PCR products were gel-purified and then used in assembly PCR to amplify the whole nsP4 gene. For the assembly PCR, 50 pmol of the outer primers
20 (3 and 4) and about 10 ng of each PCR fragment was used. The reaction volume was 100 µl, containing 4 units of Taq or Pwo polymerase, 0.1 mM dNTPs and 1.5 mM MgSO₄. The PCR conditions were as followed:

 95°C for 2 minutes, followed by 5 cycles of 92°C (45 seconds), 58°C (30
seconds), 72°C (120 seconds) and followed by 25 cycles of 92°C (45 seconds), 64°C
25 (30 seconds), 72°C (120 seconds).

 The obtained PCR fragment was purified as described above and the eluate was digested with 20 units of *SpeI* and *HpaI* in an appropriate buffer. The fragment was gel-purified and ligated into gel-purified *SpeI/HpaI* restricted SinRep5 vector. The correct sequence of the obtained vector was checked by DNA sequencing.

30 Overnight digestion of SinRep5-nsP4mut and SinRep5-nsP2mut with *SpeI/HpaI* and gel purification of the nsP4 fragment and sinRep-nsP2mut vector. The nsP4mut fragment was ligated into the SinRep5-nsP2mut vector. The final step

was cloning the nsp gene into the 987/SinRep5 vector (Bredenbeek, P. *et al.*, *J. Virol.* 67:6439-6446 (1993)) using *Cla*I and *Hpa*I as restriction endonucleases, the resulting vector was named pCytTs (SEQ ID NO:1).

5 Vector pCytTS2.1 (SEQ ID NO:2,) contains a different multiple cloning site than vector pCytTS, the RSV promoter is replaced by the CMV promoter, an SV40 intron sequence is included in the replicon and a puromycin resistance marker under control of the SV40 promoter is integrated on the plasmid. These elements can be cloned into pCytTS (SEQ ID NO: 1) in the following way:

10 Oligonucleotides Cyt-Link-FOR 5'- CTAGATTAATTAAGTCGAG GCGCGCCG-3' (SEQ ID NO: 11) and Cyt-Link-Rev 5'- GGCCCGGCGCGCCTCGAGTTAATTAAT-3' (SEQ ID NO:12) are hybridized and ligated into *Xba*I/*Bsp*120I digested vector pCytTS. This step introduces a different multiple cloning site into pCytTS (IntermediateI).

15 The CMV promoter is introduced by assembled PCR using the following oligonucleotides:

 5'CMV: 5'-ATAAGAATGCCGGCGGATCCGGCCATTAGC-3' (SEQ ID No:13)

 3'SinCMV: 5'-CCGTCAATACGGTTCACTAAACGAGCTCTG
20 CTTATATAGACC-3' (SEQ ID No: 14)

 5'CMVSin: 5'-GCTCGTTTAGTGAACCGTATTGACGGCGTA
GTACACAC-3' (SEQ ID No: 15)

 3'Sin: 5'- ACGTCGGCCTCAATTTGCGG-3' (SEQ ID No: 16)

25 The singular restriction site used for the final cloning step (*Ngo*AIV) is underlined. Two PCR reactions are carried out as described above using either oligonucleotide 5'CMV 1 (SEQ ID NO:13) and 3'SinCMV (SEQ ID NO:14) or 5'CMVSin (SEQ ID NO:15) and 3'Sin (SEQ ID NO:16). As template DNA vector pLNCX (Miller et al., 1989, *Biotechniques*, 7, 980-982) and vector pCytTS can be used, respectively. Both PCR products are gel-purified and then used in assembly
30 PCR. For the assembly PCR the outer primers (5'CMV and 3'Sin) are used, resulting in an 3089 bp size PCR fragment. The obtained PCR fragment is purified as described above and the eluate is digested with *Ngo*AIV and *Eco*47III in an

appropriate buffer. The fragment is gel-purified and ligated into gel-purified *Nga*IV/*Eco*47III restricted IntermediateI. This cloning step replaces the RSV promoter in pCytTS with the CMV promoter (IntermediateII).

5 The SV40 intron is amplified by PCR from vector pcDNA1.1amp (Invitrogen) using the following oligonucleotides:

SVIntron-FOR: 5'-GCGCGCGGGCCCAGAGGATCTTTGTGA AGG-3'
(SEQ ID NO:17)

SVIntron-REV: 5'-GCGCGCGCGCCGCTACATCAAATATTT TTCC-3'
(SEQ ID NO:18)

10 The PCR conditions can be chosen as followed: 94°C for 2 minutes, followed by 30 cycles of 94°C (45 seconds), 52°C (30 seconds), 72°C (60 seconds) and followed by a 7 min elongation step at 72°C. The obtained PCR fragment is purified using the Qia spin PCR kit (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311). Finally the purified PCR fragment is digested in the corresponding
15 buffer with *Bsp*120I/*Not*I and ligated into *Bsp*120I digested IntermediateII. The orientation of the insert is controlled by restriction enzyme analysis (IntermediateIII). The SV40 intron is thus introduced behind the multiple cloning site.

To introduce the puromycin resistance marker, vector pPUR (Clontech Laboratories, Inc., Cat. No. 6156-1) is modified first. Vector pPUR is
20 digested with *Xba*I and the linearized vector is treated with Klenow polymerase in order to get blunt ends. Religation of the vector destroys the *Xba*I site. The resulting vector serves as DNA template in the PCR amplification of the puromycin resistance marker. The following oligonucleotides are used:

5'SV40PUR: 5'- ACGTACGCGTGCGGCCGCGTTAGGGTGTGG
25 AAAGTCCCC-3' (SEQ ID NO:19)

3'SV40PUR: 5'- ACGTACGCGTTGGACAAACCACAACACTAG AATGC-3'
(SEQ ID NO:20)

The singular restriction site used for the cloning step (*Mlu*I) is underlined. PCR conditions can be chosen as described before with an annealing
30 temperature at 45°C. The resulting PCR fragment is purified as described above, digested with *Mlu*I and ligated into *Mlu*I digested IntermediateIII. The resulting

vector pCytTs2.1 (SEQ ID NO:2,) contains the puromycin resistance marker under control of the SV40 promoter.

EXAMPLE 2

5 COMBINATION OF THE TEMPERATURE SENSITIVE pCytTs SYSTEM WITH THE SELF REPLICATING EBNA SYSTEM

This system was generated in order to rapidly generate cell populations, which inducibly express a gene of interest. In this system, the tightly regulated pCytTs system is combined with the episomally replicating EBV system.

- 10 A. Construction of various pCytTs constructs containing either the EBNA origin of replication (OriP) alone or in conjunction with the replication initiation factor EBNA1.

15 In order to be able to subclone either the origin of replication of Epstein-Barr virus alone or together with the replication initiation factor EBNA1 we first subcloned this two cassettes into a shuttle vector from which they can easily be transferred into our pCytTs system with or without various inserts. Combinatorial constructs with pCytTs mentioned in the Examples such as pCytTs-OriP, pCytTs-SEAP-OPE, pCytTs-SEAP or the like generally contain a CMV promoter. The
20 sequence of the pCytTs with the CMV promoter is given in SEQ ID NO: 2 (pCytTs2.1; see Example 1). For sake of simplicity, however, the mentioned combinatorial constructs are abbreviated by simply stating pCytTs. The Epstein-Barr virus origin of replication OriP was amplified by PCR from pCEP4 (Invitrogen) using primers PH 51 and PH 52. The forward primer (PH 51) corresponds to
25 nucleotides 261-280 of pCep4 and the reverse primer (PH 52) to nucleotides 2221-2239 of pCep4. PCR was performed using pfx polymerase (BRL) according to manufacturer's recommendations. The resulting PCR product (nucleotides 261-2239 of pCep4) was then cloned into pGEMTeasy vector (Invitrogen), which had previously been digested with Eco RI and treated with Klenow polymerase in order
30 to get blunt ends. The resulting plasmid was termed pGemT-OriP. In order to get a

subclonable cassette containing OriP and EBNA1 gene, pCep4 was digested with Dra III and Cla I. The resulting 4651 bp fragment was purified and blunt ends were generated by T4 DNA polymerase treatment. This fragment was then subcloned into pGEMTeasy vector (Invitrogen), which had previously been digested with Eco RI and treated with Klenow polymerase in order to get blunt ends. The resulting construct was termed pGemT-OPE. From these plasmids either the origin of replication alone (pGemT-OriP) or together with the replication initiation factor (pGemT-OPE) could be excised by Not I digestion and transferred into Not I sites of the various CytTs vectors (*e.g.* pCytTs vectors containing the RSV or CMV promoter (SEQ ID NO: 1 or SEQ ID NO: 2) which resulted in the vectors pCytTs-OriP (SEQ ID NO: 21), and pCytTs-OPE (SEQ ID NO: 22). pCytTs-IFN β , pCytTs-EGFP, pCytTs-SEAP and pCytTs-cEPO vectors were linearized with Not I and the Not I fragment of pGemT-OPE was introduced by ligation leading to following constructs pCytTs-IFN β -OPE pCytTs-EGFP-OPE (Fig. 1), pCytTs-SEAP-OPE and pCytTs-EPO-OPE. Similarly pCytTs-EPO-OriP and pCytTs-EGFP-OriP were generated by cloning the Not I fragment from pGemT-OriP into the Not I site of pCytTs-cEPO and pCytTs-EGFP respectively.

Initial experiments were performed with a construct in which a slightly truncated form of OriP had been subcloned into pCytTs-EGFP. pCep4 (Invitrogen) was digested with Dra III and Nsi I and the 1923 nucleotide fragment (position 200-2124 from pCep4) was then treated with T4 polymerase in order to get blunt ends. The blunted fragment was then subcloned into pCytTs-EGFP, which had previously been digested with Not I and blunted with T4 DNA polymerase. The resulting plasmid was termed pCytTs-EGFP-OriPd.

B. Analysis of pCytTs-EGFP-OriPd in 293 EBNA cells

We first analyzed whether the introduction of Epstein-Barr virus derived origin of replication (OriP) into the pCytTs system would improve the inducible expression of a gene of interest on selected cell populations. We therefore transfected 293-EBNA cells (Invitrogen) with pCytTs-EGFP-OriPd. These cells constitutively

express Epstein-Barr virus replication initiation factor EBNA1 and support OriP mediated DNA replication. 293-EBNA cells were transfected with pCytTs-EGFP-OriPd using Lipofectamine 2000 (BRL) according to the manufacturer's recommendation. 36h after transfection cells were split 1 to 3 in presence of 1µg/ml puromycin. After 1 to 2 days all non-transfected cells had detached and the resistant cells were further expanded under puromycin selection for another week. In order to asses the expression from the new vector, resistant cell populations were passed to two plates of which one was kept at 37°C (uninduced) and one was shifted to 29°C for induction of EGFP expression. 24h after induction cells were harvested by trypsinisation and analysed by flow cytometry. The results are shown in Table 1. Upon induction cell populations carrying pCytTs-EGFP-OriPd were to 80% EGFP positive and displayed strong EGFP expression. In contrast cells, which had not been induced showed only weak EGFP expression and only about 8% of the cells, were green. Considering that stable cell populations containing CytTs-EGFP would only yield about 2-5% EGFP positive cells upon induction (data not shown) these results suggest that the OriP, which was introduced in the new vector, is functional and that it significantly improves the number of cells in a cell population, which expresses the gene of interest.

	% GFP positive cells	
	37°C	29°C
pCytTS-EGFP-Orip	8.2%	79%

Table 1: GFP expression with pCytTS-EGFP-OriPd 293-EBNA cells 1day after induction.

C. Comparison of pCytTs-EGFP, pCytTs-EGFP-OriP and pCytTs-EGFP-OPE in BHK cells

The following constructs, pCytTs-EGFP, pCytTs-EGFP-OriP and pCytTs-EGFP-OPE, were next examined in BHK (Baby hamster kidney) cells which are known to show tight regulation for the CytTs system. BHK cells were transfected with CytTs-EGFP, CytTs-EGFP-OriP and CytTs-EGFP-OPE using Lipofectamine 2000 (BRL) according to the manufacturer's recommendation. 24h after transfection cells were split 1 to 3 in presence of 5µg/ml puromycin. After 2 to 3 days all non-transfected cells had detached and the resistant cells were further expanded under puromycin selection for another week. Resistant cell populations were passed to two plates of which one was kept at 37 °C (non-induced) and one was shifted to 29 °C for induction of replicon replication and EGFP expression. Three days after induction cells were harvested and analysed by flow cytometry. The results are shown in Table 2. About 20% of the cell populations carrying CytTs-EGFP or CytTs-EGFP-OriP were EGFP positive upon induction, whereas four times more cells - up to 90% of the cells transfected with CytTs-EGFP-OPE did express EGFP after temperature induction. Thus, the inventive expression systems turned out to be very tight since less than 0.5% of the uninduced population showed weak EGFP expression.

	% GFP positive cells	
	37°C	29°C
pCytTS-EGFP	0.20	19
pCytTS-EGFP-OriP	0.00	21
pCytTS-EGFP-OPE	0.00	93

Table 2: GFP expression with pCytTS-EGFP, pCytTS-EGFP-OriP and pCytTS-EGFP- in BHK cells after 3 days of induction.

D. Inducible expression of pCytTs-IFNβ-OPE and pCytTs-SEAP-OPE in BHK cells

We further analyzed constructs which contain both OriP and the replication initiation factor EBNA1 (pCytTs-IFN β -OPE) with other genes of interest in BHK cells. These constructs do not depend on cell lines which have an integrated copy of the EBNA1 gene. As a gene of interest we choose IFN β , which had already been expressed with the conventional CytTs system. To test whether our new vector system with its episomal replication could result in high, inducible levels of expression on selected cell populations we transfected BHK cells with pCytTs-IFN β -OPE (*see* Example 2A) using Lipofectamine 2000 (BRL) according to the manufacturer's recommendation. 36h after transfection cells were split 1 to 3 in the presence of 5 μ g/ml puromycin. After 1 to 2 days all non-transfected cells had detached and the resistant cells were further expanded under puromycin selection for another 1 - 2 weeks. In order to assess the expression from the new vector, resistant cell populations were passed to two plates one of which was kept at 37°C (uninduced) and one was shifted to 29°C for induction of IFN β expression. Tissue culture supernatants were harvested from the uninduced control plate and 3, 4, 7 and 9 day after induction from the plate that had been shifted to 29 °C. The IFN β production at the different time points was then quantitated by ELISA (Kit of Fujirebio Inc., Tokyo, Japan). The induced cell populations reached a maximal level of 1,000,000 IU/ml after 7-9 days of induction and no IFN production could be detected in cells kept at non-permissive temperature Table 3. These results show that our new vector system combining the CytTs system with the episomally replicating EBNA system leads to high production levels in BHK cells on selected populations. The expression levels are at least as high as the ones obtained from a stable cell clone, which resulted from 4 subcloning rounds (data not shown). In addition the new system is still tightly regulated as demonstrated by up to 300,000 fold induction observed upon shifting the cells to 29°C.

Days	IU/ml	Fold induction
uninduced	3	
3	544410	159651
4	717990	210554

7	988880	289994
9	994140	291537

Table 3: IFN- β production with pCytTS-IFN-OPE in BHK cells after different days of induction.

Besides IFN β production we also analysed SEAP production with our new vector system in BHK cells. Selected cell populations were generated as described under C using construct pCytTs-SEAP-OPE. After two weeks of selection and propagation the resistant cells were passed into two plates of which one was kept at 37°C and the other was shifted to 29°C for induction. Samples were taken from the uninduced cell populations and from the induced cell population 3, 4, 7 and 9 days after induction. SEAP concentrations were measured in these samples using a kinetic ELISA for SEAP activity. The results are shown in Table 4.

Days	SEAP (mg/l)
uninduced	0
3	22
4	66
7	125
9	268

Table 4: SEAP production with pCytTS-SEAP-OPE after different days of induction

E. Comparison of the EPO production derived from pCytTs-OriP-EPO or with pCytTs-OPE-EPO in 293 EBNA cells

We further investigated whether an additional copy of the EBNA-1 gene on the same plasmid leads to higher expression of the gene of interest in 293-EBNA cells. 293-EBNA cells were transfected either with pCytTs-OriP-EPO or with pCytTS-OPE-EPO (see Example 2A) using Lipofectamin 2000 (BRL) according to the manufacturer's recommendation. One day after transfection the cells were split 1

to 2.5 in the presence of 0.8 µg/ml puromycin. A stable cell pool was obtained by further keeping these cells under selective pressure for one week. Resistant cell populations were then passed to two plates. Once the cells had attached, one plate was kept at 37°C and one plate was shifted to 29°C for induction. Cell culture supernatants were collected from both plates 4 and 8 days after induction. The erythropoietin (EPO) levels in the cell culture supernatants were then determined with an ELISA kit (R&D Systems, Minneapolis, USA). As shown in Table 5 the additional copy of the EBNA1 gene on the plasmid leads to a 15 to 20 fold increase in the EPO production compared to the plasmid which only contains the OriP. In conclusion, this experiment shows that, even in cells which contain an integrated copy, the expression of the gene of interest can be increased by an additional copy on the plasmid. These results suggest that the EBNA1 dosage may be a limiting factor for efficient replication to occur. In these experiments the EBNA1 dosage was increased by a copy on the OriP- containing plasmid. Similar results may be obtained if the EBNA1 dosage is increased in the cells either by the integration of further copies or if the additional copies would be provided on a separate plasmid in *trans*.

	EPO (mg/l)	
	pCytTS-EPO-OPE	pCytTS-EPO-OriP
uninduced	0	0.02
4	2.7	0.2
8	3.9	0.2

Table 5: Improvement of expression with additional copies of EBNA1 after 4 and 8 days of induction

EXAMPLE 3

CONSTRUCTION OF VECTORS OF THE pCytTs SYSTEM CONTAINING THE GLUTAMINE SYNTHETASE AS SELECTABLE MARKER

The glutamine synthetase gene is cloned either from hamster cells (CHO-K1, Chinese hamster ovary, ATCC, Cat. No. CCL-61; BHK21, Hamster Syrian kidney cells, DSMZ, Cat. No. ACC 61) or human cells (HEK 293 cells (ATCC, Cat. No. CRL-1573; Hela ATCC, Cat. No. CCL-2; Raji cells, ATCC, Cat. No. CCL-86; 293 EBNA , Invitrogen, Cat. No. R62007; and 143B cells, ATCC, Cat. No. CRL-8303) by the method of reverse transcription. Total RNA or cytoplasmic RNA is isolated from the cells using the RNeasy Kit (Qiagen, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311) according to the manufactures recommendation. Enrichment of poly(A)⁺ RNA from total RNA can be done by using the Oligotex mRNA Kit (Qiagen, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311). The resulting RNA can be treated with DNaseI to remove residual traces of DNA. cDNA synthesis is performed in the first step using the ThermoScript RT-PCR System (Invitrogen, Cat. No. 11146-024) with either total RNA or poly(A)⁺ -selected RNA primed with oligo(dT) or random primers according to the manufactures recommendation. Treatment of cDNA with RNase H to remove the complementary RNA prior to PCR is optional. In the second step, PCR is performed using primers specific for the hamster or human glutamine synthetase. To amplify the cDNA of hamster and human glutamine synthetase the following oligonucleotides can be used:

hamGSfor: 5'- GTCATGAAAGCTTGCCACCGCTCAGAGC ACCTTC-3'
(SEQ ID NO: 23)

ham GSrev: 5'- GACTTCTAGACTTGGGGCGGGGTGGGATGAAC-3'
(SEQ ID NO: 24)

hGSfor: 5'- GTCATGAAAGCTTTCTCGGCGACCAGAACACCTTC-3' (SEQ ID NO: 25)

hGSrev: 5'- GACTTCTAGACTTGGGGCGGGGTGGGATGAAC (SEQ ID NO: 26)

Restriction sites used for cloning are underlined (*Hind*III, *Xba*I). The resulting PCR fragment is purified using Qia spin PCR kit (Qiagen, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311).

To replace the puromycin resistance marker with the glutamine synthetase cassette in vector pCytTS2.1, the glutamine synthetase cassette is isolated from *Mlu*I

and *Bam*HI digested vector pGL3-GS. The fragment is treated with Klenow polymerase in order to get blunt ends and ligated into *Mlu*I digested vector pCytTS2.1 treated with Klenow polymerase.

5 To combine the resulting pCytTS2.1gs vector with the self replicating EBNA system either the origin of replication alone (pGEMT-OriP) or together with the replication initiation factor (pGemT-OriP) can be excised by *Not*I digestion and transferred into the *Not*I site of pCytTS2.1gs.

10 Vectors of the present invention containing the glutamine synthetase gene as selection marker are transfected in different cell lines as described above (EXAMPLE 2). Mammalian cell lines that do not express GS (glutamine synthetase) cannot survive without added glutamine in the medium. For these cell lines, a transfected GS gene can function as a selectable marker by permitting growth in a glutamine-free medium (e.g. mouse myeloma NS0 cell line). For other cell lines (e.g. CHO-K1) producing endogenous GS, methionine sulfoximine (MSX) has to be
15 added as selective agent to the glutamine-free medium.

EXAMPLE 4

20 STABILITY OF EXPRESSION IN DIFFERENT CELL LINES USING THE pCytTS-EBNA SYSTEM

Vectors of the pCytTS-EBNA system containing either the EBNA origin of replication (OriP) alone or in conjunction with the replication initiation factor
25 EBNA1 were transfected in different cell lines.

293 EBNA cells were transfected with vector pCytTS-EPO-OriP using Lipofectamine 2000 (BRL) according to the manufacturer's recommendation. One day after transfection cells were split 1 to 4 in the presence of 0.8 µg/ml puromycin. Several days after selection a stable cell pool was obtained which was further
30 expanded. To asses inducible EPO expression, resistant cell populations were trypsinized and two T25 flask were seeded with three million cells each. Once the cells had attached one T25 flask was kept at 37 °C (uninduced) and one was shifted

to 29 °C for induction of replicon replication and EPO expression. Cell culture supernatants were collected from both plates seven days after induction. The erythropoietin (EPO) level in the cell culture supernatants was determined with an ELISA kit (R&D Systems, Minneapolis, USA). A backup plate of the resistant cell pool was kept in culture and passaged about 1 to 10 times a week. To determine stability of EPO expression over several weeks, cells from the backup plate were seeded at intervals of one or two weeks again on T25 flask as described above. Cells were induced for 7 days and EPO expression was determined by ELISA (Table 6).

Weeks	EPO (mg/l)	
	37 °C	29 °C
1	< 0.1	2.3
2	< 0.1	2.8
3	< 0.1	4.4
4	< 0.1	4.9
5	< 0.1	4.5

Table 6: Stability of pCytTS-EPO-OriP in 293-EBNA cells between 1 and 5 weeks after completion of selection.

These results show that stable cell pools containing a vector system combining the pCytTS system with the episomally replicating EBNA system allow expression of EPO over several weeks without loss of productivity. EPO values in the range of 3-5 mg/l are obtained. Further on the pCytTS system is tightly regulated as at 37°C hardly no EPO production can be detected.

Having now fully described the present invention in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious to one of ordinary skill in the art that the same can be performed by modifying or changing the invention within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any specific embodiment thereof, and that such modifications or changes are intended to be encompassed within the scope of the appended claims.

All publications, patents and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains, and are herein incorporated by reference to the same extent as if each individual publication, patent or patent application was specifically and
5 individually indicated to be incorporated by reference.

CLAIMS:

- 5 1. A nucleic acid molecule comprising:
- (A) a first polynucleotide element which encodes an RNA molecule, said
 RNA molecule comprising:
- (a) at least one *cis*-acting sequence element;
- 10 (b) a first nucleotide sequence comprising a first open reading
 frame, said first open reading frame having a nucleotide
 sequence encoding an RNA-dependent RNA polymerase; and
- (c) at least one second nucleotide sequence selected from the
 group consisting of:
- (i) a second open reading frame encoding a polypeptide;
- 15 (ii) a nucleotide sequence complementary to all or a part of
 the second open reading frame of (i); and
- (iii) a nucleotide sequence encoding an untranslated RNA
 molecule or complement thereof;
- wherein said second nucleotide sequence is operably linked to
- 20 a promoter which is recognized by said RNA-dependent RNA
 polymerase;
- (B) a second polynucleotide element comprising an origin of replication;
 and
- (C) a third polynucleotide element encoding a replication initiation factor
- 25 capable of recognizing said origin of replication.
2. The nucleic acid molecule of claim 1, wherein said origin of replication is
 derived from a prokaryotic organism, a eukaryotic organism or a virus.
- 30 3. The nucleic acid molecule of claim 2, wherein said origin of replication is
 derived from a eukaryotic organism selected from the group consisting of
 yeast, mammals and insects.

4. The nucleic acid molecule of claim 1, wherein said origin of replication is derived from a DNA virus that allows for episomal replication.
5. The nucleic acid molecule of claim 1, wherein said origin of replication is derived from a DNA virus, preferably from a DNA virus selected from the group consisting of Papillomavirus, Polyomavirus, Adenovirus, and Hepadnaviruses.
6. The nucleic acid molecule of claim 1, wherein said origin of replication is derived from a Herpesvirus, and preferably wherein said origin of replication is derived from Epstein-Barr virus (EBV).
7. The nucleic acid molecule of claim 1, wherein said origin of replication is *oriP*.
8. The nucleic acid molecule of any one of the preceding claims, wherein said replication initiation factor is capable of operating as a plasmid maintenance factor.
9. The nucleic acid molecule of any one of the preceding claims, wherein said replication initiation factor is derived from a prokaryotic organism, a eukaryotic organism or a virus.
10. The nucleic acid molecule of any one of the preceding claims, wherein said replication initiation factor is derived from a DNA virus, preferably derived from a DNA virus selected from the group consisting of Herpesvirus, Papillomavirus, Polyomavirus, Adenovirus, and Hepadnavirus.
11. The nucleic acid molecule of any one of the preceding claims, wherein said replication initiation factor is derived from a Herpesvirus, and preferably wherein said replication initiation factor is derived from EBV.

12. The nucleic acid molecule of any one of the preceding claims, wherein said replication initiation factor is the EBNA-1 protein or a portion thereof.
- 5 13. The nucleic acid molecule of any one of the preceding claims, wherein said replication initiation factor and said origin of replication are derived from the same organism or the same virus.
- 10 14. The nucleic acid molecule of any one of the preceding claims, wherein said replication initiation factor and said origin of replication are derived from different organisms or viruses.
- 15 15. The nucleic acid molecule of any one of the preceding claims further comprising a fourth polynucleotide element, wherein said fourth polynucleotide element comprises a selection marker.
16. The nucleic acid molecule of claim 15, wherein said selection marker confers resistance to Puromycin.
- 20 17. The nucleic acid molecule of any one of the preceding claims, wherein said second open reading frame is in a translatable format after one or more RNA-dependent RNA replication events.
- 25 18. The nucleic acid molecule of any one of the preceding claims, wherein said RNA-dependent RNA polymerase is selected from the group consisting of: (a) a temperature-sensitive RNA-dependent RNA polymerase; (b) a non-cytopathic RNA-dependent RNA polymerase; and (c) a temperature-sensitive, non-cytopathic RNA-dependent RNA polymerase.
- 30 19. The nucleic acid molecule of any one of the preceding claims, wherein said RNA-dependent RNA polymerase is of viral origin, and preferably wherein said RNA-dependent RNA polymerase is of alphaviral origin.

20. The nucleic acid molecule of any one of the preceding claims, wherein said RNA-dependent RNA polymerase is derived from a Sindbis virus, a Semliki Forest virus or an Aura virus.
- 5
21. The nucleic acid molecule of any one of the preceding claims, wherein said RNA-dependent RNA polymerase is derived from a virus selected from the group consisting of Bebaru virus, Cabassou virus, Chikungunya virus, Easter equine encephalomyelitis virus, Fort morgan virus, Getah virus, Kyzylagach virus, Mayoaro virus, Middleburg virus, Mucambo virus, Ndumu virus, Pixuna virus, Tonate virus, Trinita virus, Una virus, Western equine encephalomyelitis virus, Whataroa virus, Venezuelan equine encephalomyelitis virus (VEE), and Ross River virus.
- 10
22. The nucleic acid molecule of claim 18, wherein said temperature-sensitive RNA-dependent RNA polymerase has replicase activity at temperatures below 34°C which is at least five fold greater than the replicase activity exhibited at 34°C or above.
- 15
23. The nucleic acid molecule of claim 18, wherein said temperature-sensitive RNA-dependent RNA polymerase has replicase activity at 34°C which is at least five fold lower than the replicase activity exhibited at 29°C.
- 20
24. The nucleic acid molecule of any one of the preceding claims, wherein said second open reading frame encodes a cytokine, a lymphokine, a tumor necrosis factor, an interferon, a toxic polypeptide, a prodrug or a converting enzyme.
- 25
25. The nucleic acid molecule of any one of the preceding claims, wherein said second nucleotide sequence encodes an untranslated RNA molecule selected
- 30

from the group consisting of an antisense RNA molecule, a tRNA molecule, a rRNA molecule, and a ribozyme.

26. An expression system comprising:

5 (A) a first polynucleotide element which encodes an RNA molecule, said RNA molecule comprising:

(a) at least one *cis*-acting sequence element;

10 (b) a first nucleotide sequence comprising a first open reading frame, said first open reading frame having a nucleotide sequence encoding an RNA-dependent RNA polymerase; and

(c) at least one second nucleotide sequence selected from the group consisting of:

(i) a second open reading frame encoding a polypeptide;

15 (ii) a nucleotide sequence complementary to all or a part of the second open reading frame of (i); and

(iii) a nucleotide sequence encoding an untranslated RNA molecule or complement thereof;

20 wherein said second nucleotide sequence is operably linked to a promoter which is recognized by said RNA-dependent RNA polymerase;

(B) a second polynucleotide element comprising an origin of replication; and

(C) a third polynucleotide element encoding a replication initiation factor capable of recognizing said origin of replication.

25

27. The expression system of claim 26, wherein said origin of replication is derived from a prokaryotic organism, a eukaryotic organism or a virus.

28. The expression system of claim 27, wherein said origin of replication is
30 derived from an eukaryotic organism selected from the group consisting of yeast, insects and mammals.

29. The expression system of claim 26, wherein said origin of replication is derived from a DNA virus that allows for episomal replication.
- 5 30. The expression system of claim 26, wherein said origin of replication is derived from a DNA virus, preferably from a DNA virus selected from the group consisting of Herpesvirus, Papillomavirus, Polyomavirus, Adenovirus, and Hepadnaviruses.
- 10 31. The expression system of claim 26, wherein said origin of replication is derived from a Herpesvirus, and preferably wherein said origin of replication is derived from Epstein-Barr virus (EBV).
32. The expression system of claim 26, wherein said origin of replication is *oriP*.
- 15 33. The expression system of any one of the preceding claims, wherein said replication initiation factor is capable of operating as plasmid maintenance factor.
- 20 34. The expression system of any one of the preceding claims, wherein said replication initiation factor is derived from a prokaryotic organism or a eukaryotic organism, or a virus.
- 25 35. The expression system of any one of the preceding claims, wherein said replication initiation factor is derived from a DNA virus, preferably derived from a DNA virus selected from the group consisting of Herpesvirus, Papillomavirus, Polyomavirus, Adenovirus, and Hepadnavirus.
- 30 36. The expression system of any one of the preceding claims, wherein said replication initiation factor is derived from a Herpesvirus, and preferably wherein said replication initiation factor is derived from EBV.

37. The expression system of any one of the preceding claims, wherein said replication initiation factor is the EBNA-1 protein, or portion thereof.
- 5 38. The expression system of any one of the preceding claims, wherein said replication initiation factor and said origin of replication are derived from the same organism or the same virus.
- 10 39. The expression system of any one of the preceding claims, wherein said replication initiation factor and said origin of replication are derived from different organisms.
- 15 40. The expression system of any one of the preceding claims, further comprising a fourth polynucleotide element, wherein said fourth polynucleotide element comprises a selection marker.
41. The expression system of claim 40, wherein said selection marker confers resistance to Puromycin.
- 20 42. The expression system of any one of the preceding claims, wherein said second open reading frame is in a translatable format after one or more RNA-dependent RNA replication events.
- 25 43. The expression system of any one of the preceding claims, wherein said first, second and third polynucleotide elements are DNA elements.
44. The expression system of any one of the preceding claims, wherein said first, second and third polynucleotide elements are each present on a separate nucleic acid molecule.
- 30 45. The expression system of any one of the preceding claims, wherein said first, second and third polynucleotide elements are all present on a single nucleic acid molecule.

46. The expression system of any one of the preceding claims, wherein said first and said second polynucleotide elements are present on a first nucleic acid molecule, and said third polynucleotide element is present on a second nucleic acid molecule.
47. The expression system of any one of the preceding claims, wherein said second nucleic acid molecule further comprises a fourth polynucleotide element capable of promoting the replication of said second nucleic acid molecule.
48. The expression system of any one of the preceding claims, wherein said second nucleic acid sequence further comprises an origin of replication.
49. The expression system of any one of the preceding claims, wherein said RNA-dependent RNA polymerase is selected from the group consisting of: (a) a temperature-sensitive RNA-dependent RNA polymerase; (b) a non-cytopathic RNA-dependent RNA polymerase; and (c) a temperature-sensitive, non-cytopathic RNA-dependent RNA polymerase.
50. The expression system of any one of the preceding claims, wherein said RNA-dependent RNA polymerase is of viral origin, and preferably wherein said RNA-dependent RNA polymerase is of alphaviral origin.
51. The expression system of any one of the preceding claims, wherein said RNA-dependent RNA polymerase is derived from a Sindbis virus, from a Semliki Forest virus or from an Aura virus.
52. The expression system of any one of the preceding claims, wherein said RNA-dependent RNA polymerase is derived from a virus selected from the group consisting of Bebaru virus, Cabassou virus, Chikungunya virus, Easter equine encephalomyelitis virus, Fort morgan virus, Getah virus, Kyzytagach

virus, Mayoaro virus, Middleburg virus, Mucambo virus, Ndumu virus, Pixuna virus, Tonate virus, Triniti virus, Una virus, Western equine encephalomyelitis virus, Whataroa virus, Venezuelan equine encephalomyelitis virus (VEE), and Ross River virus.

5

53. The expression system of claim 49, wherein said temperature-sensitive RNA-dependent RNA polymerase has replicase activity at temperatures below 34°C which is at least five fold greater than the replicase activity exhibited at 34°C or above.

10

54. The expression system of claim 49, wherein said temperature-sensitive RNA-dependent RNA polymerase has replicase activity at 34°C which is at least five fold lower than the replicase activity exhibited at 29°C.

15

55. The expression system of any one of the preceding claims, wherein said second open reading frame encodes a cytokine, a lymphokine, a tumor necrosis factor, an interferon, a toxic protein, a prodrug or a converting enzyme.

20

56. The expression system of any one of the preceding claims, wherein said second nucleotide sequence encodes an untranslated RNA molecule selected from the group consisting of an antisense RNA molecule, a tRNA molecule, a rRNA molecule, and a ribozyme.

25

57. A method of making a recombinant host cell comprising introducing the nucleic acid molecule of claim 1 into a host cell.

58. The method of claim 57, wherein said RNA-dependent RNA polymerase is selected from the group consisting of: (a) a temperature-sensitive RNA-dependent RNA polymerase; (b) a non-cytopathic RNA-dependent RNA

30

polymerase; and (c) a temperature-sensitive, non-cytopathic RNA-dependent RNA polymerase.

- 5 59. The method of claim 57, wherein said nucleic acid molecule is introduced by way of transfection.
60. A method of making a recombinant host cell comprising introducing the expression system of claim 26 into a host cell.
- 10 61. The method of claim 60, wherein said RNA-dependent RNA polymerase is selected from the group consisting of: (a) a temperature-sensitive RNA-dependent RNA polymerase; (b) a non-cytopathic RNA-dependent RNA polymerase; and (c) a temperature-sensitive, non-cytopathic RNA-dependent
- 15 RNA polymerase.
62. The method of claim 60, wherein said expression system is introduced by way of transfection.
- 20 63. A recombinant host cell produced by the method of claim 57.
64. A recombinant host cell produced by the method of claim 60.
65. A recombinant host cell comprising:
- 25 (A) a first polynucleotide element which encodes an RNA molecule, said RNA molecule comprising:
- (a) at least one *cis*-acting sequence element;
- (b) a first nucleotide sequence comprising a first open reading frame, said first open reading frame having a nucleotide
- 30 sequence encoding an RNA-dependent RNA polymerase; and
- (c) at least one second nucleotide sequence selected from the group consisting of:

- (i) a second open reading frame encoding a polypeptide;
- (ii) a nucleotide sequence complementary to all or a part of the open reading frame of (i); and
- (iii) a nucleotide sequence encoding an untranslated RNA molecule or complement thereof;

5

wherein said second nucleotide sequence is operably linked to a promoter which is recognized by said RNA-dependent RNA polymerase;

- (B) a second polynucleotide element comprising an origin of replication; and
- (C) a third polynucleotide element encoding a replication initiation factor capable of recognizing said origin of replication.

10

66. The recombinant host cell of claim 65, wherein said third polynucleotide is stably integrated into the genome of said host cell.

15

67. The recombinant host cell of claim 65, wherein said first and said second polynucleotide elements are present on a first nucleic acid molecule, and said third polynucleotide element is present on a second nucleic acid molecule.

20

68. The recombinant host cell of claim 65, wherein said first, second and third polynucleotide elements are provided on the same nucleic acid molecule.

69. The recombinant host cell of claim 65, wherein said host cell comprises more than one copy of said third polynucleotide.

25

70. The recombinant host cell of claim 69, wherein at least one copy of said third polynucleotide is stably integrated into the genome of said host cell.

30

71. The recombinant host cell of any one of the preceding claims, wherein said second open reading frame is in a translatable format after one or more RNA-dependent RNA replication events.

72. The recombinant host cell of any one of the preceding claims, wherein said first, second and third polynucleotide elements are DNA elements.
- 5 73. The recombinant host cell of any one of the preceding claims, further comprising a fourth polynucleotide comprising a selection marker, wherein said selection marker confers resistance to Puromycin
- 10 74. The recombinant host cell of any one of the preceding claims, wherein said second nucleic acid molecule further comprises a fifth polynucleotide element capable of promoting the replication of said second nucleic acid molecule.
- 15 75. The recombinant host cell of any one of the preceding claims, wherein said nucleotide sequence further comprises an origin of replication.
- 20 76. The recombinant host cell of any one of the preceding claims, wherein said RNA-dependent RNA polymerase is selected from the group consisting of: (a) a temperature-sensitive RNA-dependent RNA polymerase; (b) a non-cytopathic RNA-dependent RNA polymerase; and (c) a temperature-sensitive, non-cytopathic RNA-dependent RNA polymerase.
- 25 77. The recombinant host cell of any one of the preceding claims, wherein said recombinant host cell is a mammalian cell, and wherein preferably said mammalian cell is selected from the group consisting of a human cell, a primate cell and a rodent cell
- 30 78. A method for producing a polypeptide or untranslated RNA molecule, said method comprising:
(a) introducing the nucleic acid molecule of claim 1 into a host cell to produce a recombinant host cell; and

- (b) culturing said recombinant host cell under conditions suitable for expression of said polypeptide or untranslated RNA molecule.

5 79. The method of claim 78, further comprising recovering said polypeptide or untranslated RNA molecule.

80. A method for producing a polypeptide or untranslated RNA molecule, said method comprising:

- 10 (a) introducing the expression system of claim 26 into a host cell to produce a recombinant host cell; and
- (b) culturing said recombinant host cell under conditions suitable for expression of said polypeptide or untranslated RNA molecule.

15 81. The method of claim 80, further comprising recovering said polypeptide or untranslated RNA molecule.

82. A method for regulating the expression of a polypeptide or an untranslated RNA molecule, said method comprising:

- 20 (a) introducing the nucleic acid molecule of claim 18 into a host cell to produce a recombinant host cell;
- (b) growing said recombinant host cell under suitable culture conditions; and
- (c) changing the temperature of the recombinant host cell culture from:
 - 25 (i) a permissive temperature to a restrictive temperature, or
 - (ii) a restrictive temperature to a permissive temperature;

 wherein said polypeptide or untranslated RNA molecule is encoded by said first polynucleotide.

30 83. The method of claim 82, wherein said polypeptide is a polypeptide that is toxic to said host cell.

84. A method for regulating the expression of a polypeptide or an untranslated RNA molecule, said method comprising:

(a) introducing the expression system of claim 49 into a host cell to produce a recombinant host cell;

5 (b) growing said recombinant host cell under suitable culture conditions; and

(c) changing the temperature of the recombinant host cell culture from:

(i) a permissive temperature to a restrictive temperature, or

(ii) a restrictive temperature to a permissive temperature;

10 wherein said polypeptide or untranslated RNA molecule is encoded by said first polynucleotide.

85. The method of claim 84, wherein said polypeptide is a polypeptide that is toxic to said host cell.

GENE EXPRESSION SYSTEM

ABSTRACT OF THE DISCLOSURE

The present invention relates to compositions and methods that allow the production of polypeptides and untranslated RNA molecules in host cells. More specifically, the invention provides nucleic acid molecules, expression systems, recombinant host cells, methods and kits, which enable the production and/or isolation and/or purification of polypeptides and untranslated RNA molecules. The compositions and methods of the invention can be applied in a regulatable expression system that is transiently transfected into mammalian host cells.

Abstract


A probe assembly for an infrared medical thermometer without the necessity to put a plastic film probe cover onto the probe.

The probe assembly includes a plastic probe body (1), an elongated metal made hollow heat sink (2), which holds an infrared sensor unit (4) and conducts the heat increase of the sensor unit (4) during the temperature measurement, a hollow plastic cap (3) tightly fixed between the infrared sensor unit (4), metal heat sink (1), and the inner wall (11) of the probe body (1). Like this, the probe assembly is waterproof and/or airtight, thereby offering an washable probe which can be disinfected after a performed measurement. Therefore, the concern of cross-contamination between patients is reduced.

(Figure 1a)

-75-

FIG. 1



SEQUENCE LISTING

DT05 Rec'd PCT/PTO 18 FEB 2005

<110> Cytos Biotechnology AG
Ivanova, Lidia
Renner, Wolfgang A
Saudan, Philippe A

<120> Gene Expression System

<130> PA029WO

<150> US 60/404,928

<151> 2002-08-22

<150> US 60/432,259

<151> 2002-12-11

<160> 26

<170> PatentIn version 3.2

<210> 1

<211> 11282

<212> DNA

<213> Artificial Sequence

<220>

<223> cDNA

<400> 1

```

ctgacgcgcc ctgtagcggc gcattaagcg cggcgggtgt ggtggttacg cgcagcgtga      60
ccgctacact tgccagcgcc ctacgcgccg ctcttttcgc tttcttccct tcctttctcg      120
ccacgttcgc cggctttccc cgtcaagctc taaatcgggg gctcccttta gggttccgat      180
ttagtgcttt acggcacctc gacccccaaa aacttgatta gggtagtggt tcacgtagtg      240
ggccatcgcc ctgatagacg gtttttcgcc ctttgacggt ggagtccacg ttctttaata      300
gtggactctt gttccaaact ggaacaacac tcaaccctat ctcggtctat tcttttgatt      360
tataagggat tttgccgatt tcggcctatt ggttaaaaaa tgagctgatt taacaaaaat      420
ttaacgcgaa ttttaacaaa atattaacgc ttacaatttc cattcgccat tcaggctgcg      480
caactgttgg gaagggcgat cgggtgcgggc ctcttcgcta ttacgccagc tggcgaaagg      540
gggatgtgct gcaaggcgat taagttgggt aacgccaggg ttttcccagt cacgacgttg      600
taaaacgacg gccagtgagc gcgcaattaa ccctcactaa aggaacaaa agctggctag      660
tggatccagt cttatgcaat actctttagt tcttgcaaca tggtaacgat gagttagcaa      720
catgccttac aaggagagaa aaagcaccgt gcatgccgat tggtggaagt aagtggtac      780
gatcgtgcct tattaggaag gcaacagacg ggtctgacat ggattggacg aaccactgaa      840
ttccgcattg cagagatatt gtattttaagt gccctacctc gataccgtcg agattgacgg      900
cgtagtacac actattgaat caaacagccg accaattgca ctaccatcac aatggagaag      960
ccagtagtaa acgtagacgt agacccccag agtccgtttg tcgtgcaact gcaaaaaagc     1020

```

ttccccgcaat	ttgaggtagt	agcacagcag	gtcactccaa	atgaccatgc	taatgccaga	1080
gcatttttcgc	atctggccag	taaactaata	gagctggagg	ttcctaccac	agcgacgata	1140
ttggacatag	gcagcgcacc	ggctcgtaga	atgtttttccg	agcaccagta	tcattgtgtc	1200
tgccccatgc	gtagtccaga	agacccggac	cgcattgatga	aatacgccag	taaactggcg	1260
gaaaaagcgt	gcaagattac	aaacaagaac	ttgcatgaga	agattaagga	tctccggacc	1320
gtacttgata	cgccggatgc	tgaaacacca	tcgctctgct	ttcacaacga	tgttacctgc	1380
aacatgcgtg	ccgaatatcc	cgtcatgcag	gacgtgtata	tcaacgctcc	cggaactatc	1440
tatcatcagg	ctatgaaagg	cgtgcggacc	ctgtactgga	ttggcttcga	caccacccag	1500
ttcatgttct	cggctatggc	aggttcgtac	cctgcgtaca	acaccaactg	ggccgacgag	1560
aaagtccttg	aagcgcgtaa	catcggactt	tgacgcacaa	agctgagtga	aggtaggaca	1620
ggaaaattgt	cgataatgag	gaagaaggag	ttgaagcccg	ggtcgcgggt	ttattttctcc	1680
gtaggatcga	cactttatcc	agaacacaga	gccagcttgc	agagctggca	tcttccatcg	1740
gtgttccact	tgaatggaaa	gcagtcgtac	acttgccgct	gtgatacagt	ggtgagttgc	1800
gaaggctacg	tagtgaagaa	aatcaccatc	agtcccggga	tcacgggaga	aaccgtggga	1860
tacgcggtta	cacacaatag	cgagggcttc	ttgctatgca	aagttactga	cacagtaaaa	1920
ggagaacggg	tatcgttccc	tgtgtgcacg	tacatcccgg	ccaccatatg	cgatcagatg	1980
actggtataa	tggccacgga	tatatcacct	gacgatgcac	aaaaacttct	ggttgggctc	2040
aaccagcgaa	ttgtcattaa	cggtaggact	aacaggaaca	ccaacaccat	gcaaaattac	2100
cttctgccga	tcatagcaca	agggttcagc	aatggggcta	aggagcgcaa	ggatgatctt	2160
gataacgaga	aaatgctggg	tactagagaa	cgcaagctta	cgtatggctg	cttgtgggcg	2220
tttcgcacta	agaaagtaca	ttcgttttat	cgcccacctg	gaacgcagac	ctgcgtaaaa	2280
gtcccagcct	cttttagcgc	ttttcccatg	tcgtccgtat	ggacgacctc	tttgcccatg	2340
tcgctgaggc	agaaattgaa	actggcattg	caaccaaaga	aggaggaaaa	actgctgcag	2400
gtctcggagg	aattagtcac	ggaggccaag	gctgcttttg	aggatgctca	ggagggaagcc	2460
agagcggaga	agctccgaga	agcacttcca	ccattagtgg	cagacaaagg	catcgaggca	2520
gccgcagaag	ttgtctgcga	agtggagggg	ctccaggcgg	acatcggagc	agcattagtt	2580
gaaacccccgc	gcggtcacgt	aaggataata	cctcaagcaa	atgaccgtat	gatcggacag	2640
tatatcgttg	tctcgccaaa	ctctgtgctg	aagaatgcca	aactcgcacc	agcgaccccg	2700
ctagcagatc	aggttaagat	cataacacac	tccggaagat	caggaaggta	cgcggtcgaa	2760
ccatacgacg	ctaaagtact	gatgccagca	ggaggtgccc	taccatggcc	agaattccta	2820
gcactgagtg	agagcgccac	gttagtgtac	aacgaaagag	agtttgtgaa	ccgcaaacta	2880

taccacattg	ccatgcatgg	ccccgccaag	aatacagaag	aggagcagta	caagggttaca	2940
aaggcagagc	ttgcagaaac	agagtacgtg	tttgacgtgg	acaagaagcg	ttgcgttaag	3000
aaggaagaag	cctcaggtct	ggtcctctcg	ggagaactga	ccaaccctcc	ctatcatgag	3060
ctagctctgg	agggactgaa	gacccgacct	gcggtcccgt	acaaggtcga	aacaatagga	3120
gtgataggca	caccgggggtc	gggcaagtca	gctattatca	agtcaactgt	cacggcacga	3180
gatcttgтта	ccagcggaaa	gaaagaaaat	tgtcgcgaaa	ttgaggccga	cgtgctaaga	3240
ctgaggggta	tgcagattac	gtcgaagaca	gtagattcgg	ttatgctcaa	cggatgccac	3300
aaagccgtag	aagtgctgta	cgttgacgaa	gcgttcgcgt	gccacgcagg	agcactactt	3360
gccttgattg	ctatcgtcag	gccccgcaag	aaggtagtac	tatgcggaga	cccatgcaa	3420
tgcggattct	tcaacatgat	gcaactaaag	gtacatttca	atcacctga	aaaagacata	3480
tgcaccaaga	cattctacaa	gtatatctcc	cggcgttgca	cacagccagt	tacagctatt	3540
gtatcgacac	tgcattacga	tggaaagatg	aaaaccacga	accctgcaa	gaagaacatt	3600
gaaatcgata	ttacaggggc	cacaaagccg	aagccagggg	atatcatcct	gacatgtttc	3660
cgcggtggg	ttaagcaatt	gcaaatcgac	tatcccggac	atgaagtaat	gacagccgcg	3720
gcctcacaag	ggctaaccag	aaaaggagtg	tatgccgtcc	ggcaaaaagt	caatgaaaac	3780
ccactgtacg	cgatcacatc	agagcatgtg	aacgtgttgc	tcacccgcac	tgaggacagg	3840
ctagtgtgga	aaaccttgca	gggcgaccca	tggattaagc	agcccactaa	catacctaaa	3900
ggaaactttc	aggctactat	agaggactgg	gaagctgaac	acaagggaat	aattgctgca	3960
ataaacagcc	ccactccccg	tgccaatccg	ttcagctgca	agaccaacgt	ttgctgggcg	4020
aaagcattgg	aaccgatact	agccacggcc	ggtatcgta	ttaccggttg	ccagtggagc	4080
gaactgttcc	cacagtttgc	ggatgacaaa	ccacattcgg	ccatttacgc	cttagacgta	4140
atttgcatта	agtttttcgg	catggacttg	acaagcggac	tgttttctaa	acagagcatc	4200
ccactaacgt	accatccccg	cgattcagcg	aggccggtag	ctcattggga	caacagccca	4260
ggaacccgca	agtatgggta	cgatcacgcc	attgccgcgc	aactctcccc	tagatttccg	4320
gtgttccagc	tagctgggaa	gggcacacaa	cttgatttgc	agacggggag	aaccagagtt	4380
atctctgcac	agcataacct	ggtcccgggtg	aaccgcaatc	ttcctcacgc	cttagtcccc	4440
gagtacaagg	agaagcaacc	cggcccgggtc	aaaaaattct	tgaaccagtt	caaacaccac	4500
tcagtacttg	tggtatcaga	ggaaaaaatt	gaagctcccc	gtaagagaat	cgaatggatc	4560
gccccgattg	gcatagccgg	tgcagataag	aactacaacc	tggcttttcg	gtttccgccc	4620
caggcacggg	acgacctgg	gttcatcaac	attggaacta	aatacagaaa	ccaccacttt	4680
cagcagtgcg	aagaccatgc	ggcgacctta	aaaacccttt	cgcgttcggc	cctgaattgt	4740
ttaaactcag	gaggcacccct	cgtggtgaag	tcctatggct	acgccgaccg	caacagtga	4800

gacgtagtca	ccgctcttgc	cagaaagttt	gtcaggggtgt	ctgcagcgag	accagattgt	4860
gtctcaagca	atacagaaat	gtacctgatt	ttccgacaac	tagacaacag	ccgtacacgg	4920
caattcaccc	cgcaccatct	gaattgcgtg	atttcgtccg	tgtatgaggg	tacaagagat	4980
ggagttggag	ccgcgcctgc	ataccgcacc	aaaagggaga	atattgctga	ctgtcaagag	5040
gaagcagttg	tcaacgcagc	caatccgctg	ggtagaccag	gcgaaggagt	ctgccgtgcc	5100
atctataaac	gttggccgac	cagttttacc	gattcagcca	cggagacagg	caccgcaaga	5160
atgactgtgt	gcctaggaaa	gaaagtgatc	cacgcggtcg	gccctgattt	ccggaagcac	5220
ccagaagcag	aagccttgaa	attgctacaa	aacgcctacc	atgcagtggc	agacttagta	5280
aatgaacata	acatcaagtc	tgtcgccatt	ccactgctat	ctacaggcat	ttacgcagcc	5340
ggaaaagacc	gccttgaagt	atcacttaac	tgcttgacaa	ccgcgctaga	cagaactgac	5400
gcggacgtaa	ccatctattg	cctggataag	aagtggaagg	aaagaatcga	cgcggcactc	5460
caacttaagg	agtctgtaac	agagctgaag	gatgaagata	tggagatcga	cgatgagtta	5520
gtatggattc	atccagacag	ttgcttgaag	ggaagaaagg	gattcagtac	tacaaaagga	5580
aaattgtatt	cgtacttcga	aggcaccaaa	ttccatcaag	cagcaaaaaga	catggcggag	5640
ataaaggtcc	tgttccctaa	tgaccaggaa	agtaatgaac	aactgtgtgc	ctacatattg	5700
ggtgagacca	tggaagcaat	ccgcgaaaag	tgcccgggtcg	accataaccc	gtcgtctagc	5760
ccgccccaaa	cgttgccgtg	cctttgcatg	tatgccatga	cgccagaaaag	ggccacaga	5820
cttagaagca	ataacgtcaa	agaagttaca	gtatgctcct	ccacccccct	tcctaagcac	5880
aaaattaaga	atgttcagaa	ggttcagtgc	acgaaagtag	tcctgtttta	tccgcacact	5940
cccgcattcg	ttcccgcctg	taagtacata	gaagtgccag	aacagcctac	cgctcctcct	6000
gcacaggccg	aggaggcccc	cgaagttgta	gcgacaccgt	caccatctac	agctgataac	6060
acctcgcttg	atgtcacaga	catctcactg	gatatggatg	acagtagcga	aggctcactt	6120
ttttcgagct	ttagcggatc	ggacaactct	attactagta	tggacagttg	gtcgtcagga	6180
cctagttcac	tagagatagt	agaccgaagg	caggtgggtg	tggctgacgt	tcatgccgtc	6240
caagagcctg	cccctattcc	accgccaaag	ctaaagaaga	tggcccgcc	ggcagcggca	6300
agaaaagagc	ccactccacc	ggcaagcaat	agctctgagt	ccctccacct	ctcttttgg	6360
ggggtatcca	tgctccctcg	atcaattttc	gacggagaga	cggcccgcca	ggcagcggta	6420
caacccttg	caacaggccc	cacggatgtg	cctatgtctt	tcggatcggt	ttccgacgga	6480
gagattgatg	agctgagccg	cagagtaact	gagtccgaac	ccgtcctggt	tggatcattt	6540
gaaccggg	aagtgaactc	aattatatcg	tcccgatcag	ccgtatcttt	tccactacgc	6600
aagcagagac	gtagacgcag	gagcaggagg	actgaatact	gactaaccgg	ggtaggtggg	6660

tacatatattt	cgacggacac	aggccctggg	cacttgcaaa	agaagtcctgt	tctgcagaac	6720
cagcttacag	aaccgacctt	ggagcgcaat	gtcctggaaa	gaattcatgc	cccggtgctc	6780
gacacgtcga	aagaggaaca	actcaaactc	aggtaccaga	tgatgcccac	cgaagccaac	6840
aaaagtaggt	accagtctcg	taaagtagaa	aatcagaaaag	ccataaccac	tgagcgacta	6900
ctgtcaggac	tacgactgta	taactctgcc	acagatcagc	cagaatgcta	taagatcacc	6960
tatccgaaac	cattgtactc	cagtagcgta	ccggcgaaact	actccgatcc	acagttcgct	7020
gtagctgtct	gtaacaacta	tctgcatgag	aactatccga	cagtagcatc	ttatcagatt	7080
actgacgagt	acgatgctta	cttgगतatg	gtagacgaga	cagtcgcatg	cctggatact	7140
gcaaccttct	gccccgctaa	gcttagaagt	tacccgaaaa	aacatgagta	tagagccccg	7200
aatatccgca	gtgcggttcc	atcagcgatg	cagaacacgc	tacaaaatgt	gctcattgcc	7260
gcaactaaaa	gaaattgcaa	cgtcacgcag	atgcggtgaac	tgccaacact	ggactcagcg	7320
acattcaatg	tcgaatgctt	tcgaaaatat	gcatgtaatg	acgagtattg	ggaggagttc	7380
gctcggaaag	caattaggat	taccactgag	tttgtcaccg	catatgtagc	tagactgaaa	7440
ggccctaagg	cgcgcgcact	at ttgcaaag	acgtataatt	tggtcccatt	gcaagaagtg	7500
cctatggata	gattcgatcat	ggacatgaaa	agagacgtga	aagttacacc	aggcacgaaa	7560
cacacagaag	aaagaccgaa	agtacaagtg	atacaagccg	cagaacccct	ggcgactgct	7620
tacttatgcg	ggattcaccg	ggaattagtg	cgtaggctta	cggccgtctt	gcttccaaac	7680
attcacacgc	tttttgacat	gtcggcgag	gattttgatg	caatcatagc	agaacacttc	7740
aagcaaggcg	acccggtact	ggagacggat	atcgcatcat	tcgacaaaag	ccaagacgac	7800
gctatggcgt	taaccggtct	gatgatcttg	gaggacctgg	gtgtggatca	accactactc	7860
gacttgatcg	agtgcgcctt	tggagaaata	tcatccaccc	atctacctac	gggtactcgt	7920
tttaaattcg	gggcgatgat	gaaatccgga	atgttcctca	cactttttgt	caacacagtt	7980
ttgaatgtcg	ttatcgccag	cagagtacta	gaagagcggc	ttaaaacgtc	cagatgtgca	8040
gcgttcattg	gcgacgacaa	catcatacat	ggagtagtat	ctgacaaaaga	aatggctgag	8100
aggtgcgcca	cctgggtcaa	catggagggt	aagatcatcg	acgcagtcac	cgggtgagaga	8160
ccaccttact	tctgcggcgg	at ttatcttg	caagattcgg	ttacttccac	agcgtgccgc	8220
gtggcggatc	ccctgaaaag	gctgtttaag	ttgggtaaac	cgctcccagc	cgacgacgag	8280
caagacgaag	acagaagacg	cgctctgcta	gatgaaacaa	aggcgtgggt	tagagtaggt	8340
ataacaggca	ctttagcagt	ggccgtgacg	acccggtatg	aggtagacaa	tattacacct	8400
gtcctactgg	cattgagaac	ttttgccag	agcaaaagag	cattccaagc	catcagaggg	8460
gaaataaagc	atctctacgg	tggtcctaaa	tagtcagcat	agtacatttc	atctgactaa	8520
tactacaaca	ccaccacctc	tagacgcgta	gatctcacgt	gagcatgcag	gccttggggc	8580

caatgatccg	accagcaaaa	ctcgatgtac	ttccgaggaa	ctgatgtgca	taatgcatca	8640
ggctggtaca	ttagatcccc	gcttaccgcg	ggcaatatag	caacactaaa	aactcgatgt	8700
acttccgagg	aagcgcagtg	cataatgctg	cgcagtggtg	ccacataacc	actatattaa	8760
ccatttatct	agcggacgcc	aaaaactcaa	tgtattttctg	aggaagcgtg	gtgcataatg	8820
ccacgcagcg	tctgcataac	ttttattatt	tcttttatta	atcaacaaaa	ttttgttttt	8880
aacattttcaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaagggaa	ttcccaactt	8940
gtttatttgca	gcttataatg	gttacaataa	aagcaatagc	atcacaaatt	tcacaaataa	9000
agcatttttt	tcaactgcatt	ctagttgtgg	tttgtccaaa	ctcatcaatg	tatcttatca	9060
tgtctggatc	cgtcgagacg	cgtccaattc	gccctatagt	gagtcgtatt	acgcgcgctt	9120
ggcgtaatca	tggtcatagc	tgtttcctgt	gtgaaattgt	tatccgctca	caattccaca	9180
caacatacga	gccggaagca	taaagtgtaa	agcctggggg	gcctaataag	tgagctaact	9240
cacattaatt	gcgttgcgct	cactgcccg	tttcagctcg	ggaaacctgt	cgtgccagct	9300
gcattaatga	atcggccaac	gcgcggggag	aggcggtttg	cgtattgggc	gctcttcgc	9360
ttctcgcctc	actgactcgc	tgcgctcggt	cgttcggctg	cggcgagcgg	tatcagctca	9420
ctcaaaggcg	gtaatacggg	tatccacaga	atcaggggat	aacgcaggaa	agaacatgtg	9480
agcaaaaggc	cagcaaaagg	ccaggaaccg	taaaaaggcc	gcgttgctgg	cgtttttcca	9540
taggctccgc	ccccctgacg	agcatcacia	aaatcgacgc	tcaagtcaga	ggtggcgaaa	9600
cccgacagga	ctataaagat	accaggcggt	tccccctgga	agctccctcg	tgcgctctcc	9660
tgttccgacc	ctgccgctta	ccggatacct	gtccgccttt	ctcccttcgg	gaagcgtggc	9720
gctttctcaa	tgctcacgct	gtaggtatct	cagttcggtg	taggtcgttc	gctccaagct	9780
gggctgtgtg	cacgaacccc	ccgttcagcc	cgaccgctgc	gccttatccg	gtaactatcg	9840
tcttgagtcc	aaccgggtaa	gacacgactt	atcgccactg	gcagcagcca	ctggtaacag	9900
gattagcaga	gcgaggtatg	taggcggtgc	tacagagttc	ttgaagtggg	ggcctaacta	9960
cggctacact	agaaggacag	tatttggtat	ctgcgctctg	ctgaagccag	ttaccttcgg	10020
aaaaagagtt	ggtagctctt	gatccggcaa	acaaaccacc	gctggtagcg	gtggtttttt	10080
tgtttgcaag	cagcagatta	cgcgcagaaa	aaaaggatct	caagaagatc	ctttgatctt	10140
ttctacgggg	tctgacgctc	agtggaacga	aaactcacgt	taagggat	tggtcatgag	10200
attatcaaaa	aggatcttca	cctagatcct	tttaaattaa	aatgaagtt	ttaaatcaat	10260
ctaaagtata	tatgagtaaa	cttggtctga	cagttaccaa	tgcttaatac	gtgaggcacc	10320
tatctcagcg	atctgtctat	ttcgttcac	catagttgcc	tgactccccg	tcgtgtagat	10380
aactacgata	cgggaggggc	taccatctgg	ccccagtgct	gcaatgatac	cgcgagaccc	10440

```

acgctcaccg gctccagatt tatcagcaat aaaccagcca gccggaaggc ccgagcgcag 10500
aagtggtcct gcaactttat ccgcctccat ccagtctatt aattggtgcc gggaagctag 10560
agtaagtagt tcgccagtta atagtttgcg caacggtggt gccattgcta caggcatcgt 10620
gggtgtcacgc tcgtcgtttg gtatggcttc attcagctcc gggtcccaac gatcaaggcg 10680
agttacatga tcccccatgt tgtgcaaaaa agcgggttagc tccttcgggc ctccgatcgt 10740
tgtcagaagt aagttggccg cagtgttatc actcatgggt atggcagcac tgcataattc 10800
tcttactgtc atgccatccg taagatgctt ttctgtgact ggtgagtact caaccaagtc 10860
attctgagaa tagtgtatgc ggcgaccgag ttgctcttgc ccggcgtcaa tacgggataa 10920
taccgcgcca catagcagaa ctttaaaagt gctcatcatt ggaaaacgtt cttcggggcg 10980
aaaactctca aggatcttac cgctgttgag atccagttcg atgtaacca ctcgtgcacc 11040
caactgatct tcagcatctt ttactttcac cagcgtttct gggtgagcaa aacaggaag 11100
gcaaaatgcc gcaaaaaagg gaataagggc gacacggaaa tggtgaatac tcatactctt 11160
cctttttcaa tattattgaa gcatttatca gggttattgt ctcatgagcg gatacatatt 11220
tgaatgtatt tagaaaaata aacaaatagg gggtccgcgc acatttcccc gaaaagtgcc 11280
ac 11282

```

```

<210> 2
<211> 13068
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> cDNA

```

```

<400> 2
ctgacgcgcc ctgtagcggc gcattaagcg cggcggggtgt ggtgggttacg cgcagcgtga 60
ccgctacact tgccagcgcc ctagcgcccg ctcttttcgc tttcttccct tcctttctcg 120
ccacgttcgc cggcgggatcc ggccattagc catattatc attgggtata tagcataaat 180
caatattggc tattggccat tgcatacgtt gtatccatat cataatatgt acatttatat 240
tggctcatgt ccaacattac cgccatgttg acattgatta ttgactagtt attaatagta 300
atcaattacg gggtcattag ttcatagccc atatatggag ttccgcgtta cataacttac 360
ggtaaatggc ccgcctggct gaccgcccaa cgacccccgc ccattgacgt caataatgac 420
gtatgttccc atagtaacgc caatagggac tttccattga cgtcaatggg tggagtattt 480
acggtaaaact gccacttggc cagtacatca agtgtatcat atgccaagta cgccccctat 540
tgacgtcaat gacggtaaat ggcccgcctg gcattatgcc cagtacatga ccttatggga 600
ctttcctact tggcagtaca tctacgtatt agtcatcgct attaccatgg tgatgcgggt 660
ttggcagtag atcaatgggc gtggatagcg gtttgactca cggggatttc caagtctcca 720

```

ccccattgac	gtcaatggga	gtttgttttg	gcacccaaaat	caacgggact	ttccaaaatg	780	
tcgtaacaac	tccgccccat	tgacgcaa	gggcggtagg	cgtgtacggt	gggaggtcta	840	
tataagcaga	gctcgtttag	tgaaccgtat	tgacggcgta	gtacacacta	ttgaatcaaa	900	
cagccgacca	attgcactac	catcacaatg	gagaagccag	tagtaa	acgt agacgtagac	960	
ccccagagtc	cgtttgctgt	gcaactgcaa	aaaagcttcc	cgcaatttga	ggtagtagca	1020	
cagcaggtca	ctccaaatga	ccatgcta	at gccagagcat	tttcgcatct	ggccagtaaa	1080	
cta	atcgagc	tggaggttcc	taccacagcg	acgatcttgg	acataggcag	cgcaccggct	1140
cgtagaatgt	tttccgagca	ccagtatcat	tgtgtctgcc	ccatgcgtag	tccagaagac	1200	
ccggaccgca	tgatgaaata	tgccagtaaa	ctggcgga	aaa aagcgtgcaa	gattacaaac	1260	
aagaacttgc	atgagaagat	taaggatctc	cggaccgtac	ttgatacgcc	ggatgctgaa	1320	
acaccatcgc	tctgctttca	caacgatgtt	acctgcaaca	tgcg	tgccga atattccgtc	1380	
atgcaggacg	tgtatatcaa	cgctcccga	actatctatc	atcaggctat	gaaaggcgtg	1440	
cggaccctgt	actggattgg	cttcgacacc	accagttca	tgttctcggc	tatggcaggt	1500	
tcgtaccctg	cgtacaacac	caactgggcc	gacgagaaa	g tccctgaagc	gcgtaacatc	1560	
ggactttgca	gcacaaagct	gagtgaaggt	aggacaggaa	aattgtcgat	aatgaggaa	1620	
aaggagttga	agccccgggtc	gcgggtttat	ttctccgtag	gatcgacact	ttatccagaa	1680	
cacagagcca	gcttgcagag	ctggcatctt	ccatcggtgt	tccacttgaa	tggaagcag	1740	
tcgtacactt	gccgctgtga	tacagtgggtg	agttgcgaag	gctacgtagt	gaagaaaatc	1800	
accatcagtc	ccgggatcac	gggagaaa	acc gtgggatacg	cggttacaca	caatagcgag	1860	
ggcttcttgc	tatgcaaagt	tactgacaca	gtaaaaggag	aacgggtatc	gttccctgtg	1920	
tgcacgtaca	tcccggccac	catatgcgat	cagatgactg	gtataatggc	cacggatata	1980	
tcacctgacg	atgcacaaaa	acttctgggt	gggctcaacc	agcgaattgt	cattaacgggt	2040	
aggactaaca	ggaacaccaa	caccatgcaa	aattaccttc	tgccgatcat	agcacaaggg	2100	
ttcagcaaat	gggctaagga	gcgcaaggat	gatcttgata	acgagaaaat	gctgggtact	2160	
agagaacgca	agcttacgta	tggctgcttg	tgggcgtttc	gcactaagaa	agtacattcg	2220	
ttttatcgcc	cacctggaac	gcagacctgc	gtaaaagtcc	cagcctcttt	tagcgctttt	2280	
cccatgtcgt	ccgtatggac	gacctctttg	cccatgtcgc	tgaggcagaa	attgaaactg	2340	
gcattgcaac	caaagaagga	ggaaaaactg	ctgcaggtct	cggaggaatt	agtcattggag	2400	
gccaaggctg	cttttgagga	tgctcaggag	gaagccagag	cggagaagct	ccgagaagca	2460	
cttcaccat	tagtggcaga	caaaggcatc	gaggcagccg	cagaagttgt	ctgcgaagtg	2520	
gaggggctcc	aggcggacat	cggagcagca	ttagttgaaa	ccccgcgcgg	tcacgtaagg	2580	

ataatacctc	aagcaaatga	ccgtatgatc	ggacagtata	tcgttgtctc	gccaaactct	2640
gtgctgaaga	atgccaaact	cgcaccagcg	cacccgctag	cagatcaggt	taagatcata	2700
acacactccg	gaagatcagg	aaggtacgcg	gtcgaaccat	acgacgctaa	agtactgatg	2760
ccagcaggag	gtgccgtacc	atggccagaa	ttcctagcac	tgagtgagag	cgccacgtta	2820
gtgtacaacg	aaagagagtt	tgtgaaccgc	aaactatacc	acattgccat	gcatggcccc	2880
gccaaagaata	cagaagagga	gcagtacaag	gttacaaggg	cagagcttgc	agaaacagag	2940
tacgtgtttg	acgtggacaa	gaagcgttgc	gttaagaagg	aagaagcctc	aggtctggtc	3000
ctctcgggag	aactgaccaa	ccctccctat	catgagctag	ctctggaggg	actgaagacc	3060
cgacctgcgg	tcccgtacaa	ggtcgaaaca	ataggagtga	taggcacacc	ggggtcgggc	3120
aagtcagcta	ttatcaagtc	aactgtcacg	gcacgagatc	ttgttaccag	cggaaagaaa	3180
gaaaattgtc	gcgaaattga	ggccgacgtg	ctaagactga	ggggtatgca	gattacgtcg	3240
aagacagtag	attcggttat	gctcaacgga	tgccacaaag	ccgtagaagt	gctgtacgtt	3300
gacgaagcgt	tcgcgtgcca	cgcaggagca	ctacttgctt	tgattgctat	cgtcaggccc	3360
cgcaagaagg	tagtactatg	cggagacccc	atgcaatgcg	gattcttcaa	catgatgcaa	3420
ctaaaggtag	atttcaatca	ccctgaaaaa	gacatatgca	ccaagacatt	ctacaagtat	3480
atctcccggc	gttgcacaca	gccagttaca	gctattgtat	cgacactgca	ttacgatgga	3540
aagatgaaaa	ccacgaaccc	gtgcaagaag	aacattgaaa	tcgatattac	aggggccaca	3600
aagccgaagc	caggggatat	catcctgaca	tgtttccgcg	ggtagggtaa	gcaattgcaa	3660
atcgactatc	ccggacatga	agtaatgaca	gccgcggcct	cacaagggct	aaccagaaaa	3720
ggagtgtatg	ccgtccggca	aaaagtcaat	gaaaaccac	tgtacgcgat	cacatcagag	3780
catgtgaacg	tgttgctcac	ccgcactgag	gacaggctag	tgtggaaaac	cttgaggggc	3840
gacccatgga	ttaagcagct	cactaacata	cctaaaggaa	actttcaggc	tactatagag	3900
gactgggaag	ctgaacacaa	gggaataatt	gctgcaataa	acagccccac	tcccgtgcc	3960
aatccgttca	gctgcaagac	caacgtttgc	tgggcgaaag	cattggaacc	gatactagcc	4020
acggccggta	tcgtacttac	cggttgccag	tggagcgaac	tgttcccaca	gtttgcggat	4080
gacaaaccac	attcggccat	ttacgcctta	gacgtaattt	gcattaagtt	tttcggcatg	4140
gacttgacaa	gcggactggt	ttctaaacag	agcatcccac	taacgtacca	tccgcgcgat	4200
tcagcgaggc	cggtagctca	ttgggacaa	agcccaggaa	cccgaagta	tgggtacgat	4260
cacgccattg	ccgcogaact	ctcccgtaga	tttcgggtgt	tccagctagc	tgggaagggc	4320
acacaacttg	atttgagac	ggggagaacc	agagttatct	ctgcacagca	taacctggtc	4380
ccggtgaacc	gcaatcttcc	tcacgcctta	gtccccgagt	acaaggagaa	gcaaccggc	4440
ccggtcgaaa	aattcttgaa	ccagttcaaa	caccactcag	tacttgtggg	atcagaggaa	4500

aaaattgaag	ctccccgtaa	gagaatcgaa	tggatcgccc	cgattggcat	agccggtgca	4560
gataagaact	acaacctggc	tttcggggtt	ccgccgcagg	cacggtacga	cctggtgttc	4620
atcaacattg	gaactaaata	cagaaaccac	cacttttcagc	agtgcgaaga	ccatgcggcg	4680
accttaaaaa	cccttttcgcg	ttcggccctg	aattgtttta	actcaggagg	caccctcgtg	4740
gtgaagtcct	atggctacgc	cgaccgcaac	agtgaggacg	tagtcaccgc	tcttgccaga	4800
aagtttgtca	gggtgtctgc	agcgagacca	gattgtgtct	caagcaatac	agaaatgtac	4860
ctgattttcc	gacaactaga	caacagccgt	acacggcaat	tcaccccgca	ccatctgaat	4920
tgcgtgattt	cgtccgtgta	tgagggtaca	agagatggag	ttggagccgc	gccgtcatac	4980
cgcaccaaaa	gggagaatat	tgctgactgt	caagaggaag	cagttgtcaa	cgcagccaat	5040
ccgctgggta	gaccaggcga	aggagtctgc	cgtgccatct	ataaacgttg	gccgaccagt	5100
tttaccgatt	cagccacgga	gacaggcacc	gcaagaatga	ctgtgtgcct	aggaaagaaa	5160
gtgatccacg	cggtcggccc	tgatttcggg	aagcaccag	aagcagaagc	cttgaaattg	5220
ctacaaaacg	cctaccatgc	agtggcagac	ttagtaaatg	aacataacat	caagtctgtc	5280
gccattccac	tgctatctac	aggcatttac	gcagccggaa	aagaccgcct	tgaagtatca	5340
cttaactgct	tgacaaccgc	gctagacaga	actgacgcgg	acgtaaccat	ctattgcctg	5400
gataagaagt	ggaaggaaaag	aatcgacgcg	gcactccaac	ttaaggagtc	tgtaacagag	5460
ctgaaggatg	aagatatgga	gatcgacgat	gagttagtat	ggattcatcc	agacagttgc	5520
ttgaagggaa	gaaagggatt	cagtactaca	aaaggaaaat	tgtattcgta	cttcgaaggc	5580
accaaattcc	atcaagcagc	aaaagacatg	gcggagataa	aggtcctggt	ccctaattgac	5640
caggaaagta	atgaacaact	gtgtgcctac	atattgggtg	agaccatgga	agcaatccgc	5700
gaaaagtgcc	cggtcgacca	taacccgtcg	tctagccgcg	ccaaaacggt	gccgtgcctt	5760
tgcatgtatg	ccatgacgcc	agaaagggtc	cacagactta	gaagcaataa	cgtcaaagaa	5820
gttacagtat	gtcctccac	cccccttct	aagcacaaaa	ttaagaatgt	tcagaagggt	5880
cagtgcacga	aagtagtcct	gtttaatccg	cacactcccg	cattcgttcc	cgcccgtaa	5940
tacatagaag	tgccagaaca	gcctaccgct	cctcctgcac	aggccgagga	ggccccgaa	6000
gttgtagcga	caccgtcacc	atctacagct	gataacacct	cgcttgatgt	cacagacatc	6060
tcactggata	tggatgacag	tagcgaaggc	tcactttttt	cgagcttttag	cggatcggac	6120
aactctatta	ctagtatgga	cagttggtcg	tcaggacct	gttcactaga	gatagtagac	6180
cgaaggcagg	tggtggtggc	tgacgttcat	gccgtccaag	agcctgcccc	tattccaccg	6240
ccaaggctaa	agaagatggc	ccgcctggca	gcggcaagaa	aagagcccac	tccaccggca	6300
agcaatagct	ctgagtcctt	ccacctctct	tttggtgggg	tatccatgtc	cctcggatca	6360

atatttcgacg	gagagacggc	ccgccaggca	gcggtacaac	ccctggcaac	aggccccacg	6420
gatgtgcta	tgtctttcgg	atcgttttcc	gacggagaga	ttgatgagct	gagccgcaga	6480
gtaactgagt	ccgaaccogt	cctgttttga	tcatttgaac	cgggcgaagt	gaactcaatt	6540
atatcgcccc	gatcagccgt	atctttttcct	ctacgcaagc	agagacgtag	acgcaggagc	6600
aggaggactg	aatactgact	aaccggggta	ggtgggtaca	tatttttcgac	ggacacaggc	6660
cctggggcact	tgcaaaagaa	gtccgttctg	cagaaccagc	ttacagaacc	gaccttgag	6720
cgcaatgtcc	tggaagaat	tcatgccccg	gtgctcgaca	cgtcgaaaga	ggaacaactc	6780
aaactcaggt	accagatgat	gcccaccgaa	gccaacaaaa	gtaggtacca	gtctcgtaaa	6840
gtagaaaatc	agaaagccat	aaccactgag	cgactactgt	caggactacg	actgtataac	6900
tctgccacag	atcagccaga	atgctataag	atcacctatc	cgaaaccatt	gtactccagt	6960
agcgtaccgg	cgaactactc	cgatccacag	ttcgctgtag	ctgtctgtaa	caactatctg	7020
catgagaact	atccgacagt	agcatcttat	cagattactg	acgagtacga	tgcttacttg	7080
gatatggtag	acgagacagt	cgcatgcctg	gatactgcaa	ccttctgccc	cgctaagctt	7140
agaagttacc	cgaaaaaaca	tgagtataga	gccccgaata	tccgcagtg	ggttccatca	7200
gcgatgcaga	acacgctaca	aaatgtgctc	attgccgcaa	ctaaaagaaa	ttgcaacgtc	7260
acgcagatgc	gtgaactgcc	aacactggac	tcagcgacat	tcaatgtcga	atgctttcga	7320
aaatatgcat	gtaatgacga	gtattgggag	gagttcgctc	ggaagccaat	taggattacc	7380
actgagtttg	tcaccgcata	tgtagctaga	ctgaaaggcc	ctaaggccgc	cgactattt	7440
gcaaagacgt	ataatttgg	cccattgcaa	gaagtgccta	tggaatagatt	cgcatggac	7500
atgaaaagag	acgtgaaagt	tacaccaggc	acgaaacaca	cagaagaaag	accgaaagta	7560
caagtgatac	aagccgcaga	accctggcg	actgcttact	tatgcgggat	tcaccgggaa	7620
ttagtgcgta	ggcttacggc	cgtcttgctt	ccaaacattc	acacgctttt	tgacatgtcg	7680
gcggaggatt	ttgatgcaat	catagcagaa	cacttcaagc	aaggcgaccc	ggtactggag	7740
acggatatcg	catcattcga	caaaagccaa	gacgacgcta	tggcgttaac	cggtctgatg	7800
atcttgagg	acctgggtgt	ggatcaacca	ctactcgact	tgatcgagt	cgcttttga	7860
gaaatatcat	ccacccatct	acctacgggt	actcgtttta	aattcggggc	gatgatgaaa	7920
tccggaatgt	tcctcacact	ttttgtcaac	acagttttga	atgtcgttat	cgccagcaga	7980
gtactagaag	agcggcttaa	aacgtccaga	tgtgcagcgt	tcattggcga	cgacaacatc	8040
atacatggag	tagtatctga	caaagaaatg	gctgagaggt	gcgccacctg	gctcaacatg	8100
gagggttaaga	tcatcgacgc	agtcatcggt	gagagaccac	cttacttctg	cggcggattt	8160
atcttgcaag	attcggttac	ttccacagcg	tgccgcgtgg	cggatcccct	gaaaaggctg	8220
tttaagttgg	gtaaaccgct	cccagccgac	gacgagcaag	acgaagacag	aagacgcgct	8280

ctgctagatg	aaacaaaggc	gtggtttaga	gtaggtataa	caggcacttt	agcagtggcc	8340
gtgacgaccc	ggtatgaggt	agacaatatt	acacctgtcc	tactggcatt	gagaactttt	8400
gcccagagca	aaagagcatt	ccaagccatc	agaggggaaa	taaagcatct	ctacggtggt	8460
cctaaatagt	cagcatagta	catttcatct	gactaatacc	acaacaccac	cacctctaga	8520
ttaattaact	cgaggcgcgc	cgggccccaga	ggatctttgt	gaaggaaacct	tacttctgtg	8580
gtgtgacata	attggacaaa	ctacctacag	agatttaaag	ctctaaggta	aatataaaat	8640
ttttaagtgt	ataatgtgtt	aaactactga	ttctaattgt	ttgtgtattt	tagattccaa	8700
cctatggaac	tgatgaatgg	gagcagtggg	ggaatgcctt	taatgaggaa	aacctgtttt	8760
gctcagaaga	aatgccatct	agtgatgatg	aggctactgc	tgactctcaa	cattctactc	8820
ctccaaaaaa	gaagagaaaag	gtagaagacc	ccaaggactt	tccttcagaa	ttgctaagtt	8880
ttttgagtca	tgctgtgttt	agtaatagaa	ctcttgcttg	ctttgctatt	tacaccacaa	8940
aggaaaaagc	tgcactgcta	tacaagaaaa	ttatggaaaa	atatttgatg	tagcggccca	9000
atgatccgac	cagcaaaact	cgatgtactt	ccgaggaact	gatgtgcata	atgcatcagg	9060
ctggtacatt	agatccccgc	ttaccgcggg	caatatagca	acactaaaaa	ctcgatgtac	9120
ttccgaggaa	gcgcagtgca	taatgctgcg	cagtgttgcc	acataaccac	tatattaacc	9180
atztatctag	cggacgccaa	aaactcaatg	tatttctgag	gaagcgtggt	gcataatgcc	9240
acgcagcgtc	tgcataactt	ttattatttc	ttttattaat	caacaaaatt	ttgtttttaa	9300
catttcaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaag	ggaattccca	acttgtttat	9360
tgcagcttat	aatggttaca	aataaagcaa	tagcatcaca	aatttcacaa	ataaagcatt	9420
tttttctactg	cattctagtt	gtggtttgtc	caaactcatc	aatgtatctt	atcatgtctg	9480
gatccgtcga	gacgcgtgcg	gccgcgtagg	ggtgtggaaa	gtccccaggc	tccccagcag	9540
gcagaagtat	gcaaagcatg	catctcaatt	agtcagcaac	caggtgtgga	aagtccccag	9600
gctccccagc	aggcagaagt	atgcaaagca	tgcattctcaa	ttagtcagca	accatagtcc	9660
cgcccctaac	tccgcccatac	ccgcccctaa	ctccgcccag	ttccgcccata	tctccgcccc	9720
atggctgact	aatttttttt	atztatgcag	aggccgaggc	cgcctcggcc	tctgagctat	9780
tccagaagta	gtgaggaggc	ttttttggag	gcctaggctt	ttgcaaaaag	cttgcattgcc	9840
tgcaggtcgg	ccgccacgac	cgggtgccgc	accatcccct	gacccacgcc	cctgaccctt	9900
cacaaggaga	cgaccttcca	tgaccgagta	caagcccacg	gtgcgcctcg	ccaccgcgga	9960
cgacgtcccc	cgggcccgtac	gcaccctcgc	cgcgcggttc	gccgactacc	ccgccacgcg	10020
ccacaccgtc	gacccggacc	gccacatcga	gcgggtcacc	gagctgcaag	aactcttctt	10080
cacgcgcgtc	gggctcgaca	tcggcaaggt	gtgggtcgcg	gacgacggcg	ccgcggtggc	10140


```

ggctctggacc acgccggaga gcgtcgaagc gggggcggtg ttccgccgaga tcggcccgcg 10200
catggccgag ttgagcgggt cccggctggc cgcgcagcaa cagatggaag gcctcctggc 10260
gccgcaccgg cccaaggagc ccgcgtgggt cctggccacc gtcggcgtct cgcccacca 10320
ccagggcaag ggtctgggca gcgccgtcgt gctccccgga gtggaggcgg ccgagcgcgc 10380
cggggtgccc gccttcctgg agacctccgc gcccgcgaac ctccccttct acgagcggct 10440
cggcttcacc gtcaccgccg acgtcagagt gccgaagga ccgcgcacct ggtgcatgac 10500
ccgcaagccc ggtgcctgac gcccgcccca cgaccgcag cgcgcgaccg aaaggagcgc 10560
acgaccccat ggctccgacc gaagccaccc ggggcggccc cgcgcgaccc gcacccgccc 10620
ccgaggccca ccgactctag ctagaggatc ataatacagc ataccacatt tgtagagggt 10680
ttacttgctt taaaaaacct cccacacctc cccctgaacc tgaaacataa aatgaatgca 10740
attgttggtt ttaacttggt tattgcagct tataatgggt acaaataaag caatagcatc 10800
acaaatttca caaataaagc atttttttca ctgcattcta gttgtggttt gtccaacgcg 10860
tcggtaccag cttttgttcc ctttagtgag ggttaatttc gagcttggcg taatcatggt 10920
catagctggt tcctgtgtga aattgttacc cgctcacaat tccacacaac atacgagccg 10980
gaagcataaa gtgtaaagcc tggggtgcct aatgagttag ctaactcaca ttaattgcgt 11040
ttcgctcact gcccgctttc cagtcgggaa acctgtcgtg ccagctgcat taatgaatcg 11100
gccaacgcgc ggggagaggc ggtttgcgta ttgggcgctc ttccgcttcc tcgctcactg 11160
actcgctgcg ctcggtcggt cggctgcggc gagcgggtatc agctcactca aaggcggtaa 11220
tacggttacc cacagaatca ggggataacg caggaaagaa catgtgagca aaaggccagc 11280
aaaaggccag gaaccgtaaa aaggccgcgt tgctggcggt tttccatagg ctccgcccc 11340
ctgacgagca tcacaaaaat cgacgtcaa gtcagagggt gcgaaaccgc acaggactat 11400
aaagatacca ggcgtttccc cctggaagct ccctcgctgc ctctcctggt ccgaccctgc 11460
cgcttaccgg atacctgtcc gcctttctcc cttcgggaag cgtggcgctt tctcatagct 11520
cacgctgtag gtatctcagt tcggtgtagg tcgttcgctc caagctgggc tgtgtgcacg 11580
aaccccccg tccagccgac cgtgcgcct tatccggtaa ctatcgtctt gagtccaacc 11640
cggtaagaca cgacttatcg cactggcag cagccactgg taacaggatt agcagagcga 11700
ggatatgtag cggtgctaca gaggctctga agtggtggcc taactacggc tacactagaa 11760
gaacagtatt tggatatctg gctctgctga agccagttac cttcggaata agagttggta 11820
gctcttgatc cggcaacaa accaccgctg gtagcgggtg tttttttggt tgcaagcagc 11880
agattacgcg cagaaaaaaa ggatctcaag aagatccttt gatcttttct acggggctctg 11940
acgctcagtg gaacgaaaac tcacgttaag ggattttggt catgagatta tcaaaaagga 12000
tcttcaccta gatcctttta aattaaaaat gaagttttta atcaatctaa agtatatatg 12060

```

```

agtaaacttg gtctgacagt taccaatgct taatcagtga ggcacctatc tcagcgatct 12120
gtctatttcg ttcattccata gttgcctgac tccccgtcgt gtagataact acgatacggg 12180
agggcttacc atctggcccc agtgctgcaa tgataccgcg agacccacgc tcaccggctc 12240
cagatttatc agcaataaac cagccagccg gaagggccga gcgcagaagt ggtcctgcaa 12300
ctttatccgc ctccatccag tctattaatt gttgccggga agctagagta agtagttcgc 12360
cagttaatag tttgcgcaac gttgttgcca ttgctacagg catcgtggtg tcacgctcgt 12420
cgtttggtat ggcttcattc agctccggtt cccaacgatc aaggcgagtt acatgatccc 12480
ccatgtttgtg caaaaaagcg gttagctcct tcggctcctc gatcgttggtc agaagtaagt 12540
tggccgcagt gttatcactc atggttatgg cagcactgca taattctctt actgtcatgc 12600
catccgtaag atgcttttct gtgactgggt agtactcaac caagtcattc tgagaatagt 12660
gtatgcgggc accgagttgc tcttgcccgg cgtcaatacg ggataatacc gcgccacata 12720
gcagaacttt aaaagtgctc atcattggaa aacgttcttc ggggcgaaaa ctctcaagga 12780
tcttaccgct gttgagatcc agttcgatgt aaccactcg tgcacccaac tgatcttcag 12840
catcttttac tttcaccagc gtttctgggt gagcaaaaac aggaaggcaa aatgccgcaa 12900
aaaagggaat aaggcgaca cggaatgtt gaatactcat actcttcctt tttcaatatt 12960
attgaagcat ttatcagggt tattgtctca tgagcggata catatttgaa tgtatttaga 13020
aaaataaaca aataggggtt ccgcgcacat ttccccgaaa agtgccac 13068

```

```

<210> 3
<211> 25
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> primer

```

```

<400> 3
aacattgaaa tcgatattac agggg 25

```

```

<210> 4
<211> 20
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> primer

```

```

<400> 4
cgggttatgg tcgaccgggc 20

```

```

<210> 5
<211> 40
<212> DNA

```

<213> Artificial Sequence

<220>

<223> primer

<400> 5
gtgccctccc ctgagtttaa acaattcagg gccgaacgcg 40

<210> 6
<211> 32
<212> DNA
<213> Artificial Sequence

<220>

<223> primer

<400> 6
gaattgttta aactcaggag gcaccctcgt gg 32

<210> 7
<211> 30
<212> DNA
<213> Artificial Sequence

<220>

<223> primer

<400> 7
ggtagacgag acagtcgcat gcctggatac 30

<210> 8
<211> 30
<212> DNA
<213> Artificial Sequence

<220>

<223> primer

<400> 8
gtatccaggc atgcgactgt ctcgtctacc 30

<210> 9
<211> 25
<212> DNA
<213> Artificial Sequence

<220>

<223> primer

<400> 9
cagaccggtt aacgccatag cgtcg 25

<210> 10
<211> 25
<212> DNA
<213> Artificial Sequence

<220>

<223> primer

<400> 10
ctctattact agtatggaca gttgg 25

<210> 11
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> primer

<400> 11
ctagattaat taactcgagg cgcgccg 27

<210> 12
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> primer

<400> 12
ggcccggcgc gcctcgagtt aattaat 27

<210> 13
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> primer

<400> 13
ataagaatgc cggcggatcc ggccattagc 30

<210> 14
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> primer

<400> 14
ccgtcaatac gggtcactaa acgagctctg cttatataga cc 42

<210> 15
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> primer

<400> 15

gctcgtttag tgaaccgtat tgacggcgta gtacacac 38

<210> 16
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 16
 acgtcggcct caatttcgcg 20

<210> 17
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 17
 gcgcgcgggc ccagaggatc tttgtgaagg 30

<210> 18
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 18
 gcgcgcgcgg ccgctacatc aaatattttt cc 32

<210> 19
 <211> 39
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 19
 acgtacgcgt gcggccgcgt tagggtgtgg aaagtcccc 39

<210> 20
 <211> 33
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 20
 acgtacgcgt tggacaaacc acaactagaa tgc 33

<210> 21
 <211> 15081
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> cDNA

<400> 21
 ctgacgcgcc ctgtagcggc gcattaagcg cggcgggtgt ggtgggttacg cgcagcgtga 60
 ccgctacact tgccagcgcc ctagcgcccg ctccctttcgc tttcttccct tcctttctcg 120
 ccacgttcgc cggcgggatcc ggccattagc catattatc attgggtata tagcataaat 180
 caatattggc tattggccat tgcatacggt gtatccatat cataatatgt acatttatat 240
 tggctcatgt ccaacattac cgccatgttg acattgatta ttgactagtt attaatagta 300
 atcaattacg gggtcattag ttcatagccc atatatggag ttccgcgtta cataacttac 360
 ggtaaatggc ccgcctggct gaccgcccaa cgaccccgcc ccattgacgt caataatgac 420
 gtatgttccc atagtaacgc caatagggac tttccattga cgtcaatggg tggagtattt 480
 acggtaaaact gcccaacttg cagtacatca agtgtatcat atgccaagta cgccccctat 540
 tgacgtcaat gacggtaaat ggcccgctg gcattatgcc cagtacatga ccttatggga 600
 ctttctact tggcagtaca tctacgtatt agtcatcgct attaccatgg tgatgcggtt 660
 ttggcagtac atcaatgggc gtggatagcg gtttgactca cggggatttc caagtctcca 720
 cccattgac gtcaatggga gtttgttttg gcacaaaat caacgggact ttccaaaatg 780
 tcgtaacaac tccgccccat tgacgcaaat gggcggtagg cgtgtacggg gggaggtcta 840
 tataagcaga gctcgtttag tgaaccgtat tgacggcgta gtacacacta ttgaatcaaa 900
 cagccgacca attgcactac catcacaatg gagaagccag tagtaaactg agacgtagac 960
 cccagagtc cgtttgtcgt gcaactgcaa aaaagcttcc cgcaatttga ggtagtagca 1020
 cagcagggtca ctccaaatga ccatgcta at gccagagcat tttcgcatct ggccagtaaa 1080
 ctaatcgagc tggaggttcc taccacagcg acgatcttgg acataggcag cgcaccggct 1140
 cgtagaatgt tttccgagca ccagtatcat tgtgtctgcc ccatgcgtag tccagaagac 1200
 ccggaccgca tgatgaaata tgccagtaaa ctggcgga aagcgtgcaa gattacaaac 1260
 aagaacttgc atgagaagat taaggatctc cggaccgtac ttgatacgcc ggatgctgaa 1320
 acaccatcgc tctgctttca caacgatgtt acctgcaaca tgcgtgccga atattccgtc 1380
 atgcaggacg tgtatatcaa cgctcccga actatctatc atcaggctat gaaaggcgtg 1440
 cggaccctgt actggattgg cttcgacacc acccagttca tgttctcggc tatggcaggt 1500
 tcgtaccctg cgtacaacac caactgggcc gacgagaaag tccttgaagc gcgtaacatc 1560
 ggactttgca gcacaaagct gagtgaaggt aggacaggaa aattgtcgat aatgaggaag 1620

aaggagttga	agccccgggtc	gcggggtttat	ttctccgtag	gatcgacact	ttatccagaa	1680
cacagagcca	gcttgcagag	ctggcatctt	ccatcggtgt	tccacttgaa	tggaagcag	1740
tcgtacactt	gccgctgtga	tacagtgggtg	agttgcgaag	gctacgtagt	gaagaaaatc	1800
accatcagtc	ccgggatcac	gggagaaaacc	gtgggatacg	cggttacaca	caatagcgag	1860
ggcttcttgc	tatgcaaagt	tactgacaca	gtaaaaggag	aacgggtatc	gttccttctg	1920
tgcacgtaca	tcccggccac	catatgcat	cagatgactg	gtataatggc	cacggatata	1980
tcacctgacg	atgcacaaaa	acttctgggt	gggctcaacc	agcgaattgt	cattaacggt	2040
aggactaaca	ggaacaccaa	caccatgcaa	aattaccttc	tgccgatcat	agcacaaggg	2100
ttcagcaa	gggctaagga	gcgcaaggat	gatcttgata	acgagaaaat	gctgggtact	2160
agagaacgca	agcttacgta	tggctgcttg	tgggcgtttc	gcactaagaa	agtacattcg	2220
ttttatcgcc	cacctggaac	gcagacctgc	gtaaaagtcc	cagcctcttt	tagcgctttt	2280
cccatgtcgt	ccgtatggac	gacctctttg	cccatgtcgc	tgaggcagaa	attgaaactg	2340
gcattgcaac	caaagaagga	ggaaaaactg	ctgcaggtct	cggaggaatt	agtcattggag	2400
gccaaggctg	cttttgagga	tgctcaggag	gaagccagag	cggagaagct	ccgagaagca	2460
cttccaccat	tagtggcaga	caaaggcatc	gaggcagccg	cagaagttgt	ctgcgaagtg	2520
gaggggctcc	aggcggacat	cggagcagca	ttagttgaaa	ccccgcgcgg	tcacgtaagg	2580
ataatacctc	aagcaaatga	ccgtatgatc	ggacagtata	tcgttgtctc	gccaaactct	2640
gtgctgaaga	atgccaaaact	cgcaccagcg	cacccgctag	cagatcaggt	taagatcata	2700
acacactccg	gaagatcagg	aaggtagcgc	gtcgaaccat	acgacgctaa	agtactgatg	2760
ccagcaggag	gtgccgtacc	atggccagaa	ttcctagcac	tgagtgagag	cgccacgtta	2820
gtgtacaacg	aaagagagtt	tgtgaaccgc	aaactatacc	acattgccat	gcatggcccc	2880
gccaagaata	cagaagagga	gcagtacaag	gttacaagag	cagagcttgc	agaaacagag	2940
tacgtgtttg	acgtggacaa	gaagcgttgc	gttaagaagg	aagaagcctc	aggtctggtc	3000
ctctcgggag	aactgaccaa	ccctccctat	catgagctag	ctctggaggg	actgaagacc	3060
cgacctgcgg	tcccgtaaca	ggtcgaaaca	ataggagtga	taggcacacc	ggggtcgggc	3120
aagtcagcta	ttatcaagtc	aactgtcacg	gcacgagatc	ttgttaccag	cggaaagaaa	3180
gaaaattgtc	gcgaaattga	ggccgacgtg	ctaagactga	ggggtatgca	gattacgtcg	3240
aagacagtag	attcggttat	gctcaacgga	tgccacaaag	ccgtagaagt	gctgtacgtt	3300
gacgaagcgt	tcgcgtgcca	cgcaggagca	ctacttgcc	tgattgctat	cgtcaggccc	3360
cgcaagaagg	tagtactatg	cggagacccc	atgcaatgcg	gattcttcaa	catgatgcaa	3420
ctaaaggtag	atttcaatca	ccctgaaaaa	gacatatgca	ccaagacatt	ctacaagtat	3480
atctcccggc	gttgacacaca	gccagttaca	gctattgtat	cgacactgca	ttacgatgga	3540

aagatgaaaa	ccacgaaccc	gtgcaagaag	aacattgaaa	tcgatattac	aggggccaca	3600
aagccgaagc	caggggatat	catcctgaca	tgtttccgcg	ggtgggttaa	gcaattgcaa	3660
atcgactatc	ccggacatga	agtaatgaca	gccgcggcct	cacaagggct	aaccagaaaa	3720
ggagtgtatg	ccgtccggca	aaaagtcaat	gaaaaccac	tgtacgcgat	cacatcagag	3780
catgtgaacg	tgttgctcac	ccgcactgag	gacaggctag	tgtggaaaac	cttgcagggc	3840
gacccatgga	ttaagcagct	cactaacata	cctaaaggaa	actttcaggc	tactatagag	3900
gactgggaag	ctgaacacaa	gggaataatt	gctgcaataa	acagccccac	tccccgtgcc	3960
aatccgttca	gctgcaagac	caacgtttgc	tgggcgaaaag	cattggaacc	gatactagcc	4020
acggccggta	tcgtacttac	cggttgccag	tggagcgaac	tgttcccaca	gtttgcggat	4080
gacaaaccac	attcggccat	ttacgcctta	gacgtaattt	gcattaagtt	tttcggcatg	4140
gacttgacaa	gcggactggt	ttctaaacag	agcatcccac	taacgtacca	tcccgccgat	4200
tcagcgaggc	cggtagctca	ttgggacaac	agcccaggaa	cccgaagta	tgggtacgat	4260
cacgccattg	ccgccgaact	ctcccgtaga	tttccggtgt	tccagctagc	tgggaagggc	4320
acacaacttg	atttgcagac	ggggagaacc	agagttatct	ctgcacagca	taacctggtc	4380
ccggtgaacc	gcaatcttcc	tcacgcctta	gtccccgagt	acaaggagaa	gcaaccggc	4440
ccggtcgaaa	aattcttgaa	ccagttcaaa	caccactcag	tacttgtggt	atcagaggaa	4500
aaaattgaag	ctccccgtaa	gagaatcgaa	tggatcgccc	cgattggcat	agccggtgca	4560
gataagaact	acaacctggc	tttcgggttt	ccgccgcagg	cacggtacga	cctggtgttc	4620
atcaacattg	gaactaaata	cagaaaccac	cactttcagc	agtgcgaaga	ccatgcggcg	4680
accttaaaaa	ccctttcgcg	ttcggccctg	aattgtttta	actcaggagg	caccctcgctg	4740
gtgaagtcct	atggctacgc	cgaccgcaac	agtgaggacg	tagtcaccgc	tcttgccaga	4800
aagtttgtca	gggtgtctgc	agcgagacca	gattgtgtct	caagcaatac	agaaatgtac	4860
ctgattttcc	gacaactaga	caacagccgt	acacggcaat	tcaccccgca	ccatctgaat	4920
tgcgtgattt	cgtccgtgta	tgagggtaca	agagatggag	ttggagccgc	gccgtcatac	4980
cgcaccaaaa	gggagaatat	tgctgactgt	caagaggaag	cagttgtcaa	cgcagccaat	5040
ccgctgggta	gaccaggcga	aggagtctgc	cgtgccatct	ataaacgttg	gccgaccagt	5100
tttaccgatt	cagccacgga	gacaggcacc	gcaagaatga	ctgtgtgcct	aggaaagaaa	5160
gtgatccacg	cggtcggccc	tgatttccgg	aagcaccag	aagcagaagc	cttgaaattg	5220
ctacaaaacg	cctaccatgc	agtggcagac	ttagtaaatg	aacataacat	caagtctgtc	5280
gccattccac	tgctatctac	aggcatttac	gcagccggaa	aagaccgcct	tgaagtatca	5340
cttaactgct	tgacaaccgc	gctagacaga	actgacgcgg	acgtaaccat	ctattgcctg	5400

gataagaagt ggaaggaaag aatcgacgcg gcactccaac ttaaggagtc tgtaacagag	5460
ctgaaggatg aagatatgga gatcgacgat gagttagtat ggattcatcc agacagttgc	5520
ttgaagggaa gaaagggatt cagtactaca aaaggaaaat tgtattcgta cttcgaaggc	5580
accaaattcc atcaagcagc aaaagacatg gcgagataa aggtcctggt ccctaatagc	5640
caggaaagta atgaacaact gtgtgcctac atattgggtg agaccatgga agcaatccgc	5700
gaaaagtgcc cggtcgacca taaccctgcg tctagccgcg ccaaaacgtt gccgtgcctt	5760
tgcatgtatg ccatgacgcc agaaagggtc cacagactta gaagcaataa cgtcaaagaa	5820
gttacagtat gctcctccac cccccttctt aagcacaaaa ttaagaatgt tcagaaggtt	5880
cagtgcacga aagtagtcct gtttaatccg cacactcccg cattcgttcc cgcccgtaa	5940
tacatagaag tgccagaaca gcctaccgct cctcctgcac aggccgagga ggccccgaa	6000
gttgtagcga caccgtcacc atctacagct gataacacct cgcttgatgt cacagacatc	6060
tcactggata tggatgacag tagcgaaggc tcactttttt cgagcttttag cggatcggac	6120
aactctatta ctagtatgga cagttggtcg tcaggacctt gttcactaga gatagtagac	6180
cgaaggcagg tgggtggtggc tgacgttcat gccgtccaag agcctgcccc tattccaccg	6240
ccaaggctaa agaagatggc ccgcctggca gcggaagaa aagagccac tccaccggca	6300
agcaatagct ctgagtcctt ccacctctct tttgggtggg tatccatgtc cctcggatca	6360
atcttcgacg gagagacggc ccgccaggca gcggtacaac ccctggcaac aggccccacg	6420
gatgtgccta tgtcttttcg atcgttttcc gacggagaga ttgatgagct gagccgcaga	6480
gtaactgagt ccgaaccctg cctgtttgga tcatttgaac cgggcgaagt gaactcaatt	6540
atatcgtccc gatcagccgt atcttttctt ctacgcaagc agagacgtag acgcaggagc	6600
aggaggactg aatactgact aaccggggta ggtgggtaca tatttttcgac ggacacaggc	6660
cctgggcact tgcaaaagaa gtccgttctg cagaaccagc ttacagaacc gaccttggag	6720
cgcaatgtcc tggaaagaat tcatgccccg gtgctcgaca cgtcgaaaga ggaacaactc	6780
aaactcaggt accagatgat gccaccgaa gccaacaaaa gtaggtacca gtctcgtaaa	6840
gtagaaaatc agaaagccat aaccactgag cgactactgt caggactacg actgtataac	6900
tctgccacag atcagccaga atgctataag atcacctatc cgaaaccatt gtactccagt	6960
agcgtaccgg cgaactactc cgatccacag ttcgctgtag ctgtctgtaa caactatctg	7020
catgagaact atccgacagt agcatcttat cagattactg acgagtaoga tgcttacttg	7080
gatatggtag acgagacagt cgcatgcctg gatactgcaa ccttctgccc cgctaagctt	7140
agaagttacc cgaaaaaaca tgagtataga gccccgaata tccgcagtgc ggttccatca	7200
gcgatgcaga acacgctaca aaatgtgctc attgccgcaa ctaaaagaaa ttgcaacgtc	7260
acgcagatgc gtgaactgcc aacactggac tcagcgacat tcaatgtcga atgctttcga	7320

aaatatgcat gtaatgacga gtattgggag gagttcgctc ggaagccaat taggattacc	7380
actgagtttg tcaccgcata tgtagctaga ctgaaaggcc ctaaggccgc cgcactat	7440
gcaaagacgt ataatttggc cccattgcaa gaagtgccta tggatagatt cgtcatggac	7500
atgaaaagag acgtgaaagt tacaccaggc acgaaacaca cagaagaaag accgaaagta	7560
caagtgatac aagccgcaga acccctggcg actgcttact tatgcgggat tcaccgggaa	7620
ttagtgcgta ggcttacggc cgtcttgctt ccaaacattc acacgctttt tgacatgtcg	7680
gcgaggaggatt ttgatgcaat catagcagaa cacttcaagc aaggcgaccc ggtactggag	7740
acggatatcg catcattcga caaaagccaa gacgacgcta tggcgttaac cggctctgatg	7800
atcttggagg acctgggtgt ggatcaacca ctactcgact tgatcgagtg cgcctttgga	7860
gaaatatcat ccacccatct acctacgggt actcgtttta aattcggggc gatgatgaaa	7920
tccggaatgt tcctcacact ttttgtcaac acagttttga atgtcgttat cgccagcaga	7980
gtactagaag agcggtctaa aacgtccaga tgtgcagcgt tcattggcga cgacaacatc	8040
atacatggag tagtatctga caaagaaatg gctgagaggt gcgccacctg gctcaacatg	8100
gaggttaaga tcatcgacgc agtcatcggc gagagaccac cttacttctg cggcggat	8160
atcttgcaag attcggttac ttccacagcg tgccgcgtgg cggatcccct gaaaaggctg	8220
tttaagttgg gtaaacgcgt ccagccgac gacgagcaag acgaagacag aagacgcgt	8280
ctgctagatg aaacaaaggc gtggtttaga gtaggtataa caggcacttt agcagtggcc	8340
gtgacgaccc ggtatgaggt agacaatatt acacctgtcc tactggcatt gagaactttt	8400
gccagagca aaagagcatt ccaagccatc agaggggaaa taaagcatct ctacgggtgt	8460
cctaaatagt cagcatagta catttcatct gactaatacc acaacaccac cacctctaga	8520
ttaattaact cgaggcgcgc cgggccaga ggatctttgt gaaggaacct tacttctgtg	8580
gtgtgacata attggacaaa ctacctacag agatttaaag ctctaaggta aatataaaat	8640
ttttaagtgt ataatgtgtt aaactactga ttctaattgt ttgtgtat	8700
cctatggaac tgatgaatgg gagcagtggc ggaatgcctt taatgaggaa aacctgtttt	8760
gctcagaaga aatgccatct agtgatgatg aggctactgc tgactctcaa cattctactc	8820
ctccaaaaaa gaagagaaaag gtagaagacc ccaaggactt tccttcagaa ttgctaagtt	8880
ttttgagtca tgctgtgttt agtaatagaa ctcttgcttg ctttgctatt tacaccacaa	8940
aggaaaaagc tgcactgcta tacaagaaaa ttatggaaaa atatttgatg tagcggccca	9000
atgatccgac cagcaaaact cgatgtactt ccgaggaact gatgtgcata atgcatcagg	9060
ctggtacatt agatccccgc ttaccgcggg caatatagca aactaaaaa ctcgatgtac	9120
ttccgaggaa gcgcagtgca taatgctgcg cagtgttgcc acataaccac tatattaacc	9180

atttatctag	cggacgccaa	aaactcaatg	tattttctgag	gaagcgtggt	gcataatgcc	9240
acgcagcgtc	tgcataactt	ttattatttc	ttttattaat	caacaaaatt	ttgttttttaa	9300
catttcaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaag	ggaattccca	acttgtttat	9360
tgcagcttat	aatgggttaca	aataaagcaa	tagcatcaca	aatttcacaa	ataaagcatt	9420
tttttctactg	cattctagtt	gtggtttgtc	caaactcatc	aatgtatctt	atcatgtctg	9480
gatccgtcga	gacgcgtgcg	gccgcgggaa	ttcgatcttt	atgtgtaact	cttggctgaa	9540
gctcttacac	caatgctggg	ggacatgtac	ctcccagggg	cccaggaaga	ctacgggagg	9600
ctacaccaac	gtcaatcaga	ggggcctgtg	tagctaccga	taagcggacc	ctcaagaggg	9660
cattagcaat	agtgtttata	aggccccctt	gttaacccta	aacgggtagc	atatgcttcc	9720
cgggtagtag	tatatactat	ccagactaac	cctaattcaa	tagcatatgt	tacccaacgg	9780
gaagcatatg	ctatcgaatt	agggttagta	aaagggtcct	aaggaacagc	gatatctccc	9840
accccatgag	ctgtcacggg	tttatttaca	tggggtcagg	attccacgag	ggtagtgaac	9900
catttttagtc	acaagggcag	tggctgaaga	tcaaggagcg	ggcagtgaac	tctcctgaat	9960
cttcgcctgc	ttcttcattc	tccttcgttt	agctaataga	ataactgctg	agttgtgaac	10020
agtaagggtg	atgtgagggtg	ctcgaaaaca	aggtttcagg	tgacgcccc	agaataaaat	10080
ttggacgggg	ggttcagtg	tggcattgtg	ctatgacacc	aatataaccc	tcacaaaccc	10140
cttgggcaat	aaatactagt	gtaggaatga	aacattctga	atatctttta	caatagaaat	10200
ccatgggggtg	gggacaagcc	gtaaagactg	gatgtccatc	tcacacgaat	ttatggctat	10260
gggcaacaca	taatcctagt	gcaatatgat	actgggggtta	ttaagatgtg	tcccaggcag	10320
ggaccaagac	aggtgaacca	tgttggttaca	ctctatttgt	aacaagggga	aagagagtgg	10380
acgccgacag	cagcggactc	caactggtgt	ctctaacacc	cccgaataat	aaacgggggt	10440
ccacgccaat	ggggcccata	aacaaagaca	agtggccact	cttttttttg	aaattgtgga	10500
gtgggggac	gcgtcagccc	ccacacgccg	ccctgcgggt	ttggactgta	aaataagggt	10560
gtaataactt	ggctgattgt	aaccccgcta	accactgcgg	tcaaaccact	tgcccacaaa	10620
accactaatg	gcaccccggg	gaatacctgc	ataagtaggt	gggcggggcca	agataggggc	10680
gcgattgctg	cgatctggag	gacaaattac	acacacttgc	gcctgagcgc	caagcacagg	10740
gttggttggtc	ctcatattca	cgaggctcgt	gagagcacgg	tgggctaattg	ttgccatggg	10800
tagcatatac	tacccaaata	tctggatagc	atatgctatc	ctaactctata	tctgggtagc	10860
ataggctatc	ctaactctata	tctgggtagc	atatgctatc	ctaactctata	tctgggtagt	10920
atatgctatc	ctaatttata	tctgggtagc	ataggctatc	ctaactctata	tctgggtagc	10980
atatgctatc	ctaactctata	tctgggtagt	atatgctatc	ctaactctgta	tccgggtagc	11040
atatgctatc	ctaataagaga	ttagggtagt	atatgctatc	ctaatttata	tctgggtagc	11100

atatactacc	caaatatctg	gatagcatat	gctatcctaa	tctatatctg	ggtagcatat	11160
gctatcctaa	tctatatctg	ggtagcatag	gctatcctaa	tctatatctg	ggtagcatat	11220
gctatcctaa	tctatatctg	ggtagtatat	gctatcctaa	tttatatctg	ggtagcatag	11280
gctatcctaa	tctatatctg	ggtagcatat	gctatcctaa	tctatatctg	ggtagtatat	11340
gctatcctaa	tctgtatccg	ggtagcatat	gctatcctca	tgcataataca	gtcagcatat	11400
gatacccgagt	agtagagtg	gagtgctatc	ctttgcatat	gccgccacct	ccaaggggg	11460
cgtgaatttt	cgtgcttg	ccttttctg	catgcatcac	tagtgaattc	gcggccgct	11520
taggggtgtg	aaagtcccca	ggctccccag	caggcagaag	tatgcaaagc	atgcatctca	11580
attagtcagc	aaccagggtg	ggaaagtccc	caggctcccc	agcaggcaga	agtatgcaaa	11640
gcatgcatct	caattagtca	gcaaccatag	tcccgccct	aactccgccc	atcccgcctc	11700
taactccgcc	cagttccgcc	cattctccgc	cccatggctg	actaattttt	tttatttatg	11760
cagaggccga	ggccgcctcg	gcctctgagc	tattccagaa	gtagtgagga	ggcttttttg	11820
gaggcctagg	cttttgcaaa	aagcttgcat	gcctgcaggt	cggccgccac	gaccggtgcc	11880
gccaccatcc	cctgacccac	gcccctgacc	cctcacaagg	agacgacctt	ccatgaccga	11940
gtacaagccc	acggtgcgcc	tcgccacccg	cgacgacgtc	ccccgggccg	tacgaccctt	12000
cgccgcgcg	ttcgccgact	accccgccac	gcgccacacc	gtcgaccggg	accgccacat	12060
cgagcgggtc	accgagctgc	aagaactctt	cctcacgcgc	gtcgggctcg	acatcggcaa	12120
ggtgtgggtc	gcggacgacg	gcgccgcggt	ggcgggtctg	accacgccgg	agagcgtcga	12180
agcggggggc	gtgttcgccg	agatcggccc	gcgcatggcc	gagttgagcg	gttcccggct	12240
ggccgcgcag	caacagatgg	aaggcctcct	ggcgcgcac	cggcccaagg	agcccgctg	12300
gttcttgccc	accgtcgggc	tctcgcccga	ccaccagggc	aagggtctgg	gcagcgccgt	12360
cgtgctcccc	ggagtggagg	cggccgagcg	cgcgggggtg	cccgccttcc	tggagacctc	12420
cgcgccccgc	aacctcccct	tctacgagcg	gctcggcttc	accgtcaccg	ccgacgtcga	12480
ggtgcccga	ggaccgcgca	cctggtgcat	gaccgcgaag	cccgggtgcct	gacgcccgcc	12540
ccacgacccg	cagcgcccg	ccgaaaggag	cgcacgaccc	catgggtccg	accgaagcca	12600
cccggggcgg	ccccgccgac	cccgaccccg	cccccgaggc	ccaccgactc	tagctagagg	12660
atcataatca	gccataccac	atgtgtagag	gttttacttg	ctttaaaaaa	cctcccacac	12720
ctccccctga	acctgaaaca	taaaatgaat	gcaattgttg	ttgttaactt	gtttattgca	12780
gcttataatg	gttacaaata	aagcaatagc	atcacaaatt	tcacaaataa	agcatttttt	12840
tactgcatc	ctagttgtgg	tttgtccaac	gcgtcggtac	cagcttttgt	tccctttagt	12900
gagggttaat	ttcgagcttg	gcgtaatcat	ggtcatagct	gtttcctgtg	tgaaattggt	12960

atccgctcac	aattccacac	aacatacgag	ccggaagcat	aaagtgtaaa	gcctgggggtg	13020
cctaattgagt	gagctaactc	acattaattg	cgtttcgcctc	actgcccgcct	ttccagtcgg	13080
gaaacctgtc	gtgccagctg	cattaatgaa	tcggccaacg	cgcggggaga	ggcgggtttgc	13140
gtattgggcg	ctcttccgct	tcctcgcctca	ctgactcgct	gcgctcggtc	gttcggctgc	13200
ggcgagcggt	atcagctcac	tcaaaggcgg	taatacggtt	atccacagaa	tcaggggata	13260
acgcaggaaa	gaacatgtga	gcaaaaggcc	agcaaaaggc	caggaaccgt	aaaaaggccg	13320
cgttgctggc	gtttttccat	aggctccgcc	cccctgacga	gcatcacaaa	aatcgacgct	13380
caagtcagag	gtggcgaaac	ccgacaggac	tataaagata	ccaggcgttt	ccccctggaa	13440
gctccctcgt	gcgctctcct	gttccgaccc	tgccgcttac	cggataacctg	tccgcctttc	13500
tcccttcggg	aagcgtggcg	ctttctcata	gctcacgctg	taggtatctc	agttcggtgt	13560
aggctcgttcg	ctccaagctg	ggctgtgtgc	acgaaccccc	cgttcagccc	gaccgctgcg	13620
ccttatccgg	taactatcgt	cttgagtcca	acccggtaag	acacgactta	tcgccactgg	13680
cagcagccac	tggtaacagg	attagcagag	cgaggatatgt	aggcgggtgct	acagagttct	13740
tgaagtgggtg	gcctaactac	ggctacacta	gaagaacagt	atgttggtatc	tgcgctctgc	13800
tgaagccagt	taccttcgga	aaaagagttg	gtagctcttg	atccggcaaa	caaaccaccg	13860
ctggtagcgg	tggttttttt	gtttgcaagc	agcagattac	gcgcagaaaa	aaaggatctc	13920
aagaagatcc	tttgatcttt	tctacgggggt	ctgacgctca	gtggaacgaa	aactcacgtt	13980
aagggatttt	ggatcatgaga	ttatcaaaaa	ggatcttcac	ctagatcctt	ttaaattaaa	14040
aatgaagttt	taaatcaatc	taaagtatat	atgagtaaac	ttggtctgac	agttaccaat	14100
gcttaatcag	tgaggcacct	atctcagcga	tctgtctatt	tcgttcatcc	atagttgcct	14160
gactccccgt	cgtgtagata	actacgatac	gggaggggctt	accatctggc	cccagtgcctg	14220
caatgatacc	gcgagaccca	cgctcaccgg	ctccagattt	atcagcaata	aaccagccag	14280
ccggaagggc	cgagcgcaga	agtggctcctg	caactttatc	cgcctccatc	cagtctatta	14340
attgttgccg	ggaagctaga	gtaagtagtt	cgccagttaa	tagtttgccg	aacgttggtg	14400
ccattgctac	aggcatcgctg	gtgtcacgct	cgtcgtttgg	tatggcttca	ttcagctccg	14460
gttcccaacg	atcaaggcga	gttacatgat	cccccatggt	gtgcaaaaaa	gcggttagct	14520
ccttcgggtcc	tccgatcggt	gtcagaagta	agttggccgc	agtgttatca	ctcatggtta	14580
tggcagcact	gcataattct	cttactgtca	tgccatccgt	aagatgcttt	tctgtgactg	14640
gtgagtactc	aaccaagtca	ttctgagaat	agtgtatgcg	gcgaccgagt	tgctcttgcc	14700
cggcgtcaat	acgggataat	accgcgccac	atagcagaac	tttaaaagtg	ctcatcattg	14760
gaaaacgttc	ttcggggcga	aaactctcaa	ggatcttacc	gctggtgaga	tccagttcga	14820
tgtaaccac	tcgtgcaccc	aactgatctt	cagcatcttt	tactttcacc	agcgtttctg	14880

ggtgagcaaa aacaggaagg caaaatgccg caaaaaaggg aataagggcg acacggaaat 14940
 gttgaatact cataactcttc ctttttcaat attattgaag cattttatcag ggttattgtc 15000
 tcatgagcgg atacatatatt gaatgtattt agaaaaataa acaaataggg gttccgcgca 15060
 catttccccg aaaagtgcc a c 15081

<210> 22
 <211> 17753
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> cDNA

<400> 22
 ctgacgcgcc ctgtagcggc gcattaagcg cggcgggtgt ggtggttacg cgcagcgtga 60
 ccgctacact tgccagcgcc ctagcgcccg ctccctttcgc tttcttccct tcctttctcg 120
 ccacgttcgc cggcggatcc ggccattagc catattattc attgggtata tagcataaat 180
 caatattggc tattggccat tgcatacggt gtatccatat cataatatgt acatttatat 240
 tggctcatgt ccaacattac cgccatgttg acattgatta ttgactagtt attaatagta 300
 atcaattacg gggtcattag ttcatagcc atatatggag ttccgcgtta cataacttac 360
 ggtaaatggc ccgcctggct gaccgccaa cgaccccgcc ccattgacgt caataatgac 420
 gtatgttccc atagtaacgc caatagggac tttccattga cgtcaatggg tggagtattt 480
 acggtaaaact gcccacttgg cagtacatca agtgtatcat atgccaagta cgccccctat 540
 tgacgtcaat gacggtaaact ggccgcctg gcattatgcc cagtacatga ccttatggga 600
 ctttcctact tggcagtaca tctacgtatt agtcatcgct attaccatgg tgatgcgggt 660
 ttggcagtac atcaatgggc gtggatagcg gtttgactca cggggatttc caagtctcca 720
 cccattgac gtcaatggga gtttgttttg gcacaaaaat caacgggact ttccaaaatg 780
 tcgtaacaac tccgccccat tgacgcaaat gggcggtagg cgtgtacggg gggaggtcta 840
 tataagcaga gtcggtttag tgaaccgtat tgacggcgta gtacacacta ttgaatcaaa 900
 cagccgacca attgcactac catcacaatg gagaagccag tagtaaactg agacgtagac 960
 cccagagtc cgtttgctgt gcaactgcaa aaaagcttcc cgcaatttga ggtagtagca 1020
 cagcaggtca ctccaaatga ccatgctaatt gccagagcat tttcgcatct ggccagtaaa 1080
 ctaatcgagc tggaggttcc taccacagcg acgatcttgg acataggcag cgcaccggct 1140
 cgtagaatgt tttccgagca ccagtatcat tgtgtctgcc ccatgcgtag tccagaagac 1200
 ccggaccgca tgatgaaata tgccagtaaa ctggcggaag aagcgtgcaa gattacaaac 1260
 aagaacttgc atgagaagat taaggatctc cggaccgtac ttgatacgcc ggatgctgaa 1320

acaccatcgc	tctgctttca	caacgatgtt	acctgcaaca	tgcgtgccga	atattccgtc	1380
atgcaggacg	tgtatatcaa	cgctcccgga	actatctatc	atcaggctat	gaaaggcgtg	1440
cggaccctgt	actggattgg	cttcgacacc	accagttca	tgttctcggc	tatggcaggt	1500
tcgtaccctg	cgtacaacac	caactggggc	gacgagaaag	tccttgaagc	gcgtaacatc	1560
ggactttgca	gcacaaagct	gagtgaaggt	aggacaggaa	aattgtcgat	aatgaggaag	1620
aaggagttga	agcccgggtc	gcgggtttat	ttctccgtag	gatcgacact	ttatccagaa	1680
cacagagcca	gcttgcagag	ctggcatctt	ccatcggtgt	tcacttgaa	tggaaagcag	1740
tcgtacactt	gccgctgtga	tacagtgggt	agttgcgaag	gctacgtagt	gaagaaaatc	1800
accatcagtc	ccgggatcac	gggagaaacc	gtgggatacg	cggttacaca	caatagcgag	1860
ggcttcttgc	tatgcaaagt	tactgacaca	gtaaaaggag	aacgggtatc	gttccctgtg	1920
tgcacgtaca	tcccggccac	catatgcgat	cagatgactg	gtataatggc	cacggatata	1980
tcacctgacg	atgcacaaaa	acttctgggt	gggctcaacc	agcgaattgt	cattaacggt	2040
aggactaaca	ggaacaccaa	caccatgcaa	aattaccttc	tgccgatcat	agcacaaggg	2100
ttcagcaaat	gggctaagga	gcgcaaggat	gatcttgata	acgagaaaat	gctgggtact	2160
agagaacgca	agcttacgta	tggctgcttg	tgggcgtttc	gcactaagaa	agtacattcg	2220
ttttatcgcc	cacctggaac	gcagacctgc	gtaaaagtcc	cagcctcttt	tagcgctttt	2280
cccatgtcgt	ccgtatggac	gacctctttg	cccatgtcgc	tgaggcagaa	attgaaactg	2340
gcattgcaac	caaagaagga	ggaaaaactg	ctgcaggctc	cggaggaatt	agtcatggag	2400
gccaaggctg	cttttgagga	tgctcaggag	gaagccagag	cggagaagct	ccgagaagca	2460
cttccaccat	tagtggcaga	caaaggcatc	gaggcagccg	cagaagttgt	ctgcgaagtg	2520
gaggggctcc	aggcggacat	cggagcagca	ttagttgaaa	ccccgcgcgg	tcacgtaagg	2580
ataatacctc	aagcaaatga	ccgtatgatc	ggacagtata	tcgttgtctc	gccaaactct	2640
gtgctgaaga	atgccaaact	cgcaccagcg	cacccgctag	cagatcaggt	taagatcata	2700
acacactccg	gaagatcagg	aaggtagcgc	gtcgaaccat	acgacgctaa	agtactgatg	2760
ccagcaggag	gtgccgtacc	atggccagaa	ttcctagcac	tgagtgagag	cgccacgtta	2820
gtgtacaacg	aaagagagtt	tgtgaaccgc	aaactatacc	acattgccat	gcatggcccc	2880
gccaagaata	cagaagagga	gcagtacaag	gttacaaagg	cagagcttgc	agaaacagag	2940
tacgtgtttg	acgtggacaa	gaagcgttgc	gttaagaagg	aagaagcctc	aggtctggtc	3000
ctctcgggag	aactgaccaa	ccctccctat	catgagctag	ctctggaggg	actgaagacc	3060
cgacctgcgg	tcccgtacaa	ggtcgaaaca	ataggagtga	taggcacacc	ggggtcgggc	3120
aagtcagcta	ttatcaagtc	aactgtcacg	gcacgagatc	ttgttaccag	cggaagaaa	3180
gaaaattgtc	gcgaaattga	ggccgacgtg	ctaagactga	ggggtatgca	gattacgtcg	3240

aagacagtag attcggttat gctcaacgga tgccacaaag ccgtagaagt gctgtacgtt	3300
gacgaagcgt tcgcgtgcca cgcaggagca ctacttgctt tgattgctat cgtcaggccc	3360
cgcaagaagg tagtactatg cggagacccc atgcaatgcg gattcttcaa catgatgcaa	3420
ctaaagggtac atttcaatca ccctgaaaaa gacatatgca ccaagacatt ctacaagtat	3480
atctcccggc gttgcacaca gccagttaca gctattgtat cgacactgca ttacgatgga	3540
aagatgaaaa ccacgaaccc gtgcaagaag aacattgaaa tcgatattac agggggccaca	3600
aagccgaagc caggggatat catcctgaca tgtttccgcg ggtgggttaa gcaattgcaa	3660
atcgactatc ccggacatga agtaatgaca gccgcggcct cacaagggct aaccagaaaa	3720
ggagtgtatg ccgtccggca aaaagtcaat gaaaaccac tgtacgcgat cacatcagag	3780
catgtgaacg tgttgctcac ccgcactgag gacaggctag tgtggaaaac cttgcagggc	3840
gacccatgga ttaagcagct cactaacata cctaaaggaa actttcaggc tactatagag	3900
gactgggaag ctgaacacaa gggaataatt gctgcaataa acagccccac tccccgtgcc	3960
aatccgttca gctgcaagac caacgtttgc tgggcgaaag cattggaacc gatactagcc	4020
acggccggta tcgtacttac cggttgccag tggagcgaac tgttcccaca gtttgccgat	4080
gacaaaccac attcggccat ttacgcctta gacgtaattt gcattaagtt tttcggcatg	4140
gacttgacaa gcggactgtt ttctaaacag agcatcccac taacgtacca tcccgccgat	4200
tcagcgaggc cggtagctca ttgggacaac agcccaggaa cccgcaagta tgggtacgat	4260
cacgccattg ccgccgaact ctcccgtaga tttccggtgt tccagctagc tgggaagggc	4320
acacaacttg atttgagac ggggagaacc agagttatct ctgcacagca taacctggtc	4380
ccggtgaacc gcaatcttcc tcacgcctta gtccccgagt acaaggagaa gcaaccggc	4440
ccggtcgaaa aattcttgaa ccagttcaaa caccactcag tacttggtgt atcagaggaa	4500
aaaattgaag ctccccgtaa gagaatcgaa tggatcgccc cgattggcat agccggtgca	4560
gataagaact acaacctggc tttcgggttt ccgccgcagg cacggtacga cctggtgttc	4620
atcaacattg gaactaaata cagaaaccac cactttcagc agtgcaaga ccatgcggcg	4680
accttaaaaa ccctttcgcg ttccggccctg aattgtttta actcaggagg caccctcgtg	4740
gtgaagtcct atggctacgc cgaccgcaac agtgaggacg tagtcaccgc tcttgccaga	4800
aagtttgtca ggggtgtctgc agcgagacca gattgtgtct caagcaatac agaaatgtac	4860
ctgattttcc gacaactaga caacagccgt acacggcaat tcaccccgc ccatctgaat	4920
tgcgtagattt cgtccgtgta tgagggtaca agagatggag ttggagccgc gccgtcatac	4980
cgcaccaaaa gggagaatat tgctgactgt caagaggaag cagttgtcaa cgagccaat	5040
ccgctgggta gaccaggcga aggagtctgc cgtgccatct ataaacgttg gccgaccagt	5100

tttaccgatt	cagccacgga	gacaggcacc	gcaagaatga	ctgtgtgcct	aggaaagaaa	5160
gtgatccacg	cggtcggccc	tgatttccgg	aagcaccag	aagcagaagc	cttgaaattg	5220
ctacaaaacg	cctaccatgc	agtggcagac	ttagtaaatg	aacataacat	caagtctgtc	5280
gccattccac	tgtatcttac	aggcatttac	gcagccgga	aagaccgcct	tgaagtatca	5340
cttaactgct	tgacaaccgc	gctagacaga	actgacgcgg	acgtaaccat	ctattgcctg	5400
gataagaagt	ggaaggaaag	aatcgacgcg	gcactccaac	ttaaggagtc	tgtaacagag	5460
ctgaaggatg	aagatatgga	gatcgacgat	gagttagtat	ggattcatcc	agacagttgc	5520
ttgaagggaa	gaaagggatt	cagtactaca	aaaggaaaat	tgtattcgta	cttcgaaggc	5580
accaaattcc	atcaagcagc	aaaagacatg	gcggagataa	aggtcctggt	ccctaattgac	5640
caggaaagta	atgaacaact	gtgtgcctac	atattgggtg	agaccatgga	agcaatccgc	5700
gaaaagtgcc	cggtcgacca	taaccgcgtc	tctagccgc	ccaaaacggt	gccgtgcctt	5760
tgcatgtatg	ccatgacgcc	agaaagggtc	cacagactta	gaagcaataa	cgtcaaagaa	5820
gttacagtat	gtctctccac	cccccttct	aagcacaaaa	ttaagaatgt	tcagaaggtt	5880
cagtgcacga	aagtagtcct	gtttaatccg	cacactcccg	cattcgttcc	cgcccgtaa	5940
tacatagaag	tgccagaaca	gcctaccgct	cctctgcac	aggccgagga	ggccccgaa	6000
gttgtagcga	caccgtcacc	atctacagct	gataacacct	cgcttgatgt	cacagacatc	6060
tcactggata	tggatgacag	tagcgaaggc	tcactttttt	cgagcttttag	cggatcggac	6120
aactctatta	ctagtatgga	cagttggctg	tcaggacct	gttcactaga	gatagtagac	6180
cgaaggcagg	tggtggtggc	tgacgttcat	gccgtccaag	agcctgcccc	tattccaccg	6240
ccaaggctaa	agaagatggc	ccgcctggca	gcggcaagaa	aagagcccac	tccaccggca	6300
agcaatagct	ctgagtcctt	ccacctctct	tttggtgggg	tatccatgtc	cctcggatca	6360
atthttcgacg	gagagacggc	ccgccaggca	gcggtacaac	ccctggcaac	aggccccacg	6420
gatgtgccta	tgtcttttcg	atcgthttcc	gacggagaga	ttgatgagct	gagccgcaga	6480
gtaactgagt	ccgaaccctg	cctgttttga	tcatttgaac	cgggcgaagt	gaactcaatt	6540
atatcgthcc	gatcagccgt	atctthttct	ctacgcaagc	agagacgtag	acgcaggagc	6600
aggaggactg	aatactgact	aaccggggta	ggtgggtaca	tattthtcgac	ggacacaggc	6660
cctgggcact	tgcaaaagaa	gtccgttctg	cagaaccagc	ttacagaacc	gaccttggag	6720
cgcaatgtcc	tggaaagaat	tcatgccccg	gtgctcgaca	cgtcgaaaga	ggaacaactc	6780
aaactcaggt	accagatgat	gcccaccgaa	gccaacaaaa	gtaggtacca	gtctcgtaaa	6840
gtagaaaatc	agaaagccat	aaccactgag	cgactactgt	caggactacg	actgtataac	6900
tctgccacag	atcagccaga	atgctataag	atcacctatc	cgaaaccatt	gtactccagt	6960
agcgtaccgg	cgaactactc	cgatccacag	ttcgctgtag	ctgtctgtaa	caactatctg	7020

catgagaact atccgacagt agcatcttat cagattactg acgagtacga tgcttacttg	7080
gatatggtag acgagacagt cgcattgctg gatactgcaa ccttctgccc cgctaagctt	7140
agaagttacc cgaaaaaaca tgagtataga gccccgaata tccgcagtgc gggtccatca	7200
gcgatgcaga acacgctaca aaatgtgctc attgccgcaa ctaaaagaaa ttgcaacgtc	7260
acgcagatgc gtgaactgcc aacactggac tcagcgacat tcaatgtcga atgctttcga	7320
aaatatgcat gtaatgacga gtattgggag gagttcgctc ggaagccaat taggattacc	7380
actgagtttg tcaccgcata tgtagctaga ctgaaaggcc ctaaggccgc cgactattt	7440
gcaaagacgt ataatttggt cccattgcaa gaagtgccta tggatagatt cgtcatggac	7500
atgaaaagag acgtgaaagt tacaccaggc acgaaacaca cagaagaaag accgaaagta	7560
caagtgatac aagccgcaga acccctggcg actgcttact tatgcgggat tcaccgggaa	7620
ttagtgcgta ggcttacggc cgtcttgctt ccaaaccattc acacgctttt tgacatgtcg	7680
gcgaggatt ttgatgcaat catagcagaa cacttcaagc aaggcgaccc ggtactggag	7740
acggatatcg catcattcga caaaagccaa gacgacgcta tggcgtaac cggctctgatg	7800
atcttgagg acctgggtgt ggatcaacca ctactcgact tgatcgagtg cgcctttgga	7860
gaaatatcat ccacccatct acctacgggt actcgtttta aattcggggc gatgatgaaa	7920
tccggaatgt tcctcacact ttttgtcaac acagttttga atgtcgttat cgccagcaga	7980
gtactagaag agcggcttaa aacgtccaga tgtgcagcgt tcattggcga cgacaacatc	8040
atacatggag tagtatctga caaagaaatg gctgagaggt gcgccacctg gctcaacatg	8100
gaggttaaga tcatcgacgc agtcacgggt gagagaccac cttacttctg cggcggattt	8160
atcttgcaag attcgggttac ttccacagcg tgccgcgtgg cggatccccct gaaaaggctg	8220
tttaagttgg gtaaaccgct cccagccgac gacgagcaag acgaagacag aagacgcgct	8280
ctgctagatg aaacaaaggc gtggtttaga gtaggtataa caggcacttt agcagtggcc	8340
gtgacgaccc ggtatgaggt agacaatatt acacctgtcc tactggcatt gagaactttt	8400
gccagagca aaagagcatt ccaagccatc agaggggaaa taaagcatct ctacggtggt	8460
cctaaatagt cagcatagta catctcatct gactaatacc acaacaccac cacctctaga	8520
tttaattaact cgaggcgcgc cgggccaga ggatctttgt gaaggaacct tacttctgtg	8580
gtgtgacata attggacaaa ctacctacag agatttaaag ctctaaggta aatataaaat	8640
ttttaagtgt ataatgtgtt aaactactga ttctaattgt ttgtgtattt tagattccaa	8700
cctatggaac tgatgaatgg gagcagtggg ggaatgcctt taatgaggaa aacctgtttt	8760
gctcagaaga aatgccatct agtgatgatg aggctactgc tgactctcaa cattctactc	8820
ctccaaaaaa gaagagaaag gtagaagacc ccaaggactt tccttcagaa ttgctaagtt	8880

ttttgagtca	tgctgtgttt	agtaatagaa	ctcttgcttg	ctttgctatt	tacaccacaa	8940
aggaaaaagc	tgcactgcta	tacaagaaaa	ttatggaaaa	atatttgatg	tagcggccca	9000
atgatccgac	cagcaaaact	cgatgtactt	ccgaggaact	gatgtgcata	atgcatcagg	9060
ctggtacatt	agatccccgc	ttaccgcggg	caatatagca	acactaaaaa	ctcgatgtac	9120
ttccgaggaa	gcgcagtgca	taatgctgcg	cagtgttgcc	acataaccac	tatattaacc	9180
atztatctag	cggacgccaa	aaactcaatg	tatttctgag	gaagcgtggg	gcataatgcc	9240
acgcagcgtc	tgcataaactt	ttattatttc	ttttattaat	caacaaaatt	ttgtttttaa	9300
catttcaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaag	ggaattccca	acttgtttat	9360
tgcagcttat	aatggttaca	aataaagcaa	tagcatcaca	aatttcacaa	ataaagcatt	9420
tttttcaactg	cattctagtt	gtggtttgtc	caaactcatc	aatgtatctt	atcatgtctg	9480
gatccgtcga	gacgcgtgcg	gccgcgggaa	ttcgatgtgg	ctttcatcct	ggagcagact	9540
ttgcagtctg	tggactgcaa	cacaacattg	cctttatgtg	taactcttgg	ctgaagctct	9600
tacaccaatg	ctggggggaca	tgtacctccc	aggggcccag	gaagactacg	ggaggctaca	9660
ccaacgtcaa	tcagagggggc	ctgtgtagct	accgataagc	ggaccctcaa	gagggcatta	9720
gcaatagtgt	ttataaggcc	cccttggtta	ccctaaacgg	gtagcatatg	cttcccgggt	9780
agtagtatat	actatccaga	ctaaccctaa	ttcaatagca	tatgttacct	aacgggaagc	9840
atatgctatc	gaattagggg	tagtaaaagg	gtcctaagga	acagcgatat	ctcccacccc	9900
atgagctgtc	acggttttat	ttacatgggg	tcaggattcc	acgagggtag	tgaaccattt	9960
tagtcacaag	ggcagtggct	gaagatcaag	gagcgggcag	tgaactctcc	tgaatcttcg	10020
cctgcttctt	cattctcctt	cgtttagcta	atagaataac	tgctgagttg	tgaacagtaa	10080
gggtgtatgtg	agggtgctga	aaacaagggt	tcagggtgacg	cccccagaat	aaaatttgga	10140
cgggggggttc	agtgggtggca	ttgtgctatg	acaccaatat	aaccctcaca	aacccttggg	10200
gcaataaata	ctagtgtagg	aatgaaacat	tctgaatatc	tttaacaata	gaaatccatg	10260
gggtgggggac	aagccgtaaa	gactggatgt	ccatctcaca	cgaatttatg	gctatgggca	10320
acacataatc	ctagtgcaat	atgatactgg	ggttattaag	atgtgtccca	ggcagggacc	10380
aagacaggtg	aaccatgttg	ttacactcta	tttgtaacaa	ggggaaagag	agtggacgcc	10440
gacagcagcg	gactccactg	gttgtctcta	acacccccga	aaattaaacg	gggctccacg	10500
ccaatggggc	ccataaacia	agacaagtgg	ccactctttt	ttttgaaatt	gtggagtggg	10560
ggcacgcgtc	agccccacaa	cgccgccttg	cggtttttgg	ctgtaaaata	agggtgtaat	10620
aacttggtcg	attgtaacct	cgctaaccac	tgcggtcaaa	ccacttgccc	acaaaaccac	10680
taatggcacc	ccgggggaata	cctgcataag	taggtggggc	ggccaagata	ggggcgcgat	10740
tgctgcgata	tggaggacaa	attacacaca	cttgcgcttg	agcgccaagc	acagggttgt	10800

tggctcctcat attcacgagg tcgctgagag cacggtgggc taatgttgcc atgggtagca 10860
 tatactaccc aaatatctgg atagcatatg ctatcctaatt ctatatctgg gtagcatagg 10920
 ctatcctaatt ctatatctgg gtagcatatg ctatcctaatt ctatatctgg gtagtatatg 10980
 ctatcctaatt ttatatctgg gtagcatagg ctatcctaatt ctatatctgg gtagcatatg 11040
 ctatcctaatt ctatatctgg gtagtatatg ctatcctaatt ctgtatccgg gtagcatatg 11100
 ctatcctaatt agagattagg gtagtatatg ctatcctaatt ttatatctgg gtagcatata 11160
 ctacccaaat atctggatag catatgctat cctaattctat atctgggtag catatgctat 11220
 cctaattctat atctgggtag cataggctat cctaattctat atctgggtag catatgctat 11280
 cctaattctat atctgggtag tatatgctat cctaatttctat atctgggtag cataggctat 11340
 cctaattctat atctgggtag catatgctat cctaattctat atctgggtag tatatgctat 11400
 cctaattctgt atccgggtag catatgctat cctcatgcat atacagtcag catatgatac 11460
 ccagtagtag agtgggagtg ctatcctttg catatgccgc cacctcccaa gggggcgtga 11520
 attttcgctg cttgtccttt tcctgcatgc tggttgctcc cattcttagg tgaatttaag 11580
 gaggccaggc taaagccgtc gcatgtctga ttgctcacca ggtaaagtgc gctaattgtt 11640
 tccaacgcga gaaggtgttg agcgcggagc tgagtgcgtg gacaacatgg gtatgcccaa 11700
 ttgccccatg ttggggaggac gaaaatgggtg acaagacaga tggccagaaa tacaccaaca 11760
 gcacgcatga tgtctactgg ggattttattc tttagtgcgg gggaatacac ggcttttaatt 11820
 acgattgagg gcgtctccta acaagttaca tcactcctgc ccttcctcac cctcatctcc 11880
 atcacctcct tcactctcgt catctccgtc atcacctcct gcggcagccc cttccaccat 11940
 aggtggaaac caggaggaggc aatctactcc atcgtcaaag ctgcacacag tcaccctgat 12000
 attgcaggta ggagcgggct ttgtcataac aaggtcctta atcgcatcct tcaaaacctc 12060
 agcaaatata tgagtttgta aaaagaccat gaaataacag acaatggact cccttagcgg 12120
 gccaggttgt gggccggggtc cagggggccat tccaaagggg agacgactca atgggtgtaag 12180
 acgacattgt ggaatagcaa gggcagttcc tcgccttagg ttgtaaaggg aggtcttact 12240
 acctccatat acgaacacac cggcgaccca agttccttcg tcggtagtcc tttctacgtg 12300
 actcctagcc aggagagctc ttaaaccttc tgcaatgttc tcaaatttcg ggttggaacc 12360
 tccttgacca cgatgctttc caaaccaccc tccttttttg cgctgcctc catcacctg 12420
 accccggggg ccagtgcttg ggccttctcc tgggtcatct gcggggccct gctctatcgc 12480
 tcccgggggc acgtcaggct caccatctgg gccaccttct tgggtggtatt caaaataatc 12540
 ggcttcccct acagggtgga aaaatggcct tctacctgga gggggcctgc gcggtggaga 12600
 cccggatgat gatgactgac tactgggact cctgggcctc ttttctccac gtccacgacc 12660

tctccccctg	gtctttttac	gaattccccc	cctgggtctt	tcacgtcctc	taccccggcg	12720
gcctccacta	cctcctcgac	cccggcctcc	actacctcct	cgaccccggc	ctccactgcc	12780
tcctcgaccc	cggcctccac	ctcctgctcc	tgccccctct	gtccttgccc	ctcctcctgc	12840
tcctgcccct	cctgcccctc	ctgctcctgc	ccctcctgcc	cctcctgctc	ctgcccctcc	12900
tgccccctct	gtccttgccc	ctcctgcccc	tcctcctgct	cctgcccctc	ctgcccctcc	12960
tcctgctcct	gccccctcctg	ccccctcctgc	tcctgcccct	cctgcccctc	ctgctcctgc	13020
ccctcctgcc	cctcctgctc	ctgcccctcc	tgctcctgcc	cctcctgctc	ctgcccctcc	13080
tgctcctgcc	cctcctgccc	ctcctgcccc	tcctcctgct	cctgcccctc	ctgctcctgc	13140
ccctcctgcc	cctcctgccc	ctcctgctcc	tgccccctct	cctgctcctg	ccccctcctgc	13200
ccctcctgcc	cctcctcctg	ctcctgcccc	tcctgcccct	cctcctgctc	ctgcccctcc	13260
tcctgctcct	gccccctcctg	ccccctcctgc	ccctcctcct	gtccttgccc	ctcctgcccc	13320
tcctcctgct	cctgcccctc	ctcctgctcc	tgccccctct	gccccctcctg	ccccctcctcc	13380
tgctcctgcc	cctcctcctg	ctcctgcccc	tcctgcccct	cctgcccctc	ctgcccctcc	13440
tcctgctcct	gccccctcctc	ctgctcctgc	ccctcctgct	cctgcccctc	ccgtcctcgc	13500
tcctgctcct	gttccacogt	gggtcccttt	gcagccaatg	caacttggac	gtttttgggg	13560
tctccggaca	ccatctctat	gtcttgggcc	tgatcctgag	ccgccggggg	ctcctgggtct	13620
tcgcctcct	cgctcctgctc	ctcttccccg	tcctcgctcca	tggttatcac	ccccctctct	13680
ttgaggtcca	ctgcgcgcgg	agccttctgg	tccagatgtg	tctcccttct	ctcctaggcc	13740
atttccaggt	cctgtacctg	gccccctgctc	agacatgatt	cacactaaaa	gagatcaata	13800
gacatcttta	ttagacgacg	ctcagtgaat	acagggagtg	cagactcctg	ccccctccaa	13860
cagccccccc	accctcatcc	ccttcatggt	cgctgtcaga	cagatccagg	tctgaaaatt	13920
ccccatcctc	cgaaccatcc	togtctcat	caccaattac	tgcagccccg	gaaaactccc	13980
gctgaacatc	ctcaagattt	gcgtcctgag	cctcaagcca	ggcctcaaat	tcctcgctcc	14040
cctttttgct	ggacggtagg	gatggggatt	ctcgggaccc	ctcctcttcc	tcttcaaggt	14100
caccagacag	agatgctact	ggggcaacgg	aagaaaagct	gggtgcggcc	tgtgaggatc	14160
agcttatatc	actagtgaat	tgcgggccgc	gtaggggtgt	ggaaagtccc	caggctcccc	14220
agcaggcaga	agtatgcaaa	gcatgcatct	caattagtca	gcaaccaggt	gtggaaagtc	14280
cccaggctcc	ccagcaggca	gaagtatgca	aagcatgcat	ctcaattagt	cagcaaccat	14340
agtcccgcgc	ctaactccgc	ccatcccgc	cctaactccg	cccagttccg	ccattctcc	14400
gccccatggc	tgactaattt	tttttattta	tgcagaggcc	gaggccgct	cggcctctga	14460
gctattccag	aagtagtgag	gaggcttttt	tggaggccta	ggcttttgca	aaaagcttgc	14520
atgcctgcag	gtcggccgcc	acgaccggtg	ccgccaccat	cccctgaccc	acgcccctga	14580

```

ccccacaaa ggagacgacc ttccatgacc gagtacaagc ccacggtgcg cctcgccacc 14640
cgcgacgacg tcccccgggc cgtacgcacc ctgcgcgcgc cggttcgcca ctaccccgcc 14700
acgcgccaca ccgtcgaccc ggaccgccac atcgagcggg tcaccgagct gcaagaactc 14760
ttcctcacgc gcgtcgggct cgacatcggc aagggtgtggg tcgcggacga cggcgccgcg 14820
gtggcggtct ggaccacgcc ggagagcgtc gaagcggggg cggtgttcgc cgagatcggc 14880
ccgcgcatgg ccgagttgag cggttcccgc ctggccgcgc agcaacagat ggaaggcctc 14940
ctggcgccgc accggcccaa ggagcccgcg tggttcctgg ccaccgtcgc cgtctcgccc 15000
gaccaccagg gcaaggttct gggcagcgcc gtcgtgctcc ccggagtgga ggcgcccgag 15060
cgcgccgggg tgcccgcctt cctggagacc tccgcgcccc gcaacctccc cttctacgag 15120
cggctcggct tcaccgtcac cgccgacgtc gaggtgcccg aaggaccgcg cacctggtgc 15180
atgaccgcga agcccgggtg ctgacgcccg cccacgacc cgcagcgccc gaccgaaagg 15240
agcgacgac cccatggctc cgaccgaagc caccggggc ggccccgcgc accccgcacc 15300
cgcccccgag gccaccgac tctagctaga ggatcataat cagccatacc acattttag 15360
aggttttact tgctttaaaa aacctccac acctccccct gaacctgaaa cataaaatga 15420
atgcaattgt tgttgtaac ttgtttattg cagcttataa tggttacaaa taaagcaata 15480
gcatcacaaa tttcacaaat aaagcatttt tttactgca ttctagttgt ggtttgtcca 15540
acgcgtcggg accagctttt gttcccttta gtgagggtta atttcgagct tggcgtaatc 15600
atggtcatag ctgtttcctg tgtgaaattg ttatccgctc acaattccac acaacatacg 15660
agccggaagc ataaagtgt aagcctgggg tgcctaatga gtgagctaac tcacattaat 15720
tgcgtttcgc tactgcccgc ctttcagtc gggaaacctg tcgtgccagc tgcattaatg 15780
aatcggccaa cgcgcgggga gaggcggttt gcgtattggg cgctcttcgc cttcctcgct 15840
cactgactcg ctgcgctcgc tcgttcggct gcggcgagcg gtatcagctc actcaaaggc 15900
ggtaatacgg ttatccacag aatcagggga taacgcagga aagaacatgt gagcaaaagg 15960
ccagcaaaag gccaggaacc gtaaaaaggc cgcgttgctg gcgtttttcc ataggctccg 16020
ccccctgac gagcatcaca aaaatcgacg ctcaagtcag aggtggcgaa acccgacagg 16080
actataaaga taccaggcgt tttcccctgg aagctccctc gtgcgctctc ctgttccgac 16140
cctgccgctt accggatacc tgtccgcctt tctcccttcg ggaagcgtgg cgctttctca 16200
tagctcacgc tgtaggtatc tcagttcggg gtaggtcggt cgctccaagc tgggctgtgt 16260
gcacgaaccc cccgttcagc ccgaccgctg cgccttatcc ggtaactatc gtcttgagtc 16320
caaccgggta agacacgact tatcgccact ggcagcagcc actggtaaca ggattagcag 16380
agcgaggtat gtaggcgggtg ctacagagtt cttgaagtgg tggcctaact acggctacac 16440

```

tagaagaaca gtatattgga tctgcgctct gctgaagcca gttaccttcg gaaaaagagt 16500
 tggtagctct tgatccggca aacaaaccac cgctggtagc ggtgggtttt ttgtttgcaa 16560
 gcagcagatt acgcgcagaa aaaaaggatc tcaagaagat cctttgatct tttctacggg 16620
 gtctgacgct cagtggaaacg aaaactcacg ttaagggatt ttggtcatga gattatcaaa 16680
 aaggatcttc acctagatcc ttttaaatta aaaatgaagt tttaaatcaa tctaaagtat 16740
 atatgagtaa acttggtctg acagttacca atgcttaatc agtgaggcac ctatctcagc 16800
 gatctgtcta tttcgttcat ccatagtgtg ctgactcccc gtcgtgtaga taactacgat 16860
 acgggagggc ttaccatctg gccccagtgc tgcaatgata ccgcgagacc cacgctcacc 16920
 ggctccagat ttatcagcaa taaaccagcc agccggaagg gccgagcgca gaagtgggtcc 16980
 tgcaacttta tccgcctcca tccagtctat taattgttgc cgggaagcta gagtaagtag 17040
 ttcgccagtt aatagtttgc gcaacgttgt tgccattgct acaggcatcg tgggtgtcacg 17100
 ctgctggttt ggtatggctt cattcagctc cggttcccaa cgatcaaggc gagttacatg 17160
 atcccccatg ttgtgcaaaa aagcggttag ctcttcggt cctccgatcg ttgtcagaag 17220
 taagttggcc gcagtgttat cactcatggt tatggcagca ctgcataatt ctcttactgt 17280
 catgccatcc gtaagatgct tttctgtgac tggtgagtac tcaaccaagt cattctgaga 17340
 atagtgtatg cggcgaccga gttgctcttg cccggcgctca atacgggata ataccgcgcc 17400
 acatagcaga actttaaaag tgctcatcat tggaacacgt tcttcggggc gaaaactctc 17460
 aaggatctta ccgctgttga gatccagttc gatgtaaccc actcgtgcac ccaactgatc 17520
 ttcagcatct tttactttca ccagcgtttc tgggtgagca aaaacaggaa ggcaaaatgc 17580
 cgcaaaaaag ggaataaggc cgacacggaa atgttgaata ctcatactct tcctttttca 17640
 atattattga agcatttatc agggttattg tctcatgagc ggatacatat ttgaatgtat 17700
 ttagaaaaat aaacaaatag gggttccgcg cacatttccc cgaaaagtgc cac 17753

<210> 23
 <211> 33
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 23
 gtcatgaagc ttgccaccgc tcagagcacc ttc

33

<210> 24
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>

<223> primer

<400> 24

gacttctaga ctggggcggg gtgggatgaa c

31

<210> 25

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> primer

<400> 25

gtcatgaagc ttctcggcga ccagaacacc ttc

33

<210> 26

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> primer

<400> 26

gacttctaga ctggggcggg gtgggatgaa c

31

1

32